

A BASIS OF THE FIXED POINT SUBGROUP OF AN AUTOMORPHISM OF A FREE GROUP

OLEG BOGOPOLSKI* AND OLGA MASLAKOVA†

ABSTRACT. We prove that for any automorphism α of a free group F of finite rank, one can compute a basis of the fixed point subgroup $\text{Fix}(\alpha)$.

CONTENTS

1. Introduction	2
2. Preliminaries	4
3. Outline of the proof	5
4. A relative train track for an automorphism of a free group of finite rank	12
5. Graphs D_f and $CoRe(C_f)$ for a homotopy equivalence $f : \Gamma \rightarrow \Gamma$	19
6. Cancellation points in the product of two paths	28
7. Non-deletable cancellation points in exponentially growing paths	30
8. Stable r -paths and their A -decompositions	38
9. Subdivided relative train track	40
10. r -perfect and A -perfect paths	42
11. Some technical propositions	43
12. Infinite μ -subgraphs and f -orbits of some vertices of D_f	49
13. Good vertices in infinite μ -graphs	51
14. E -perfect paths	57
15. Finiteness and membership problems for μ -subgraphs	60
References	63

*Mathematisches Institut, Düsseldorf University, Düsseldorf, Germany
and Sobolev Institute of Mathematics of SB RAS, Novosibirsk, Russia.
e-mail: bogopolski@math.uni-duesseldorf.de.

†Sobolev Institute of Mathematics of SB RAS, Novosibirsk, Russia.
e-mail: o.s.maslakova@gmail.com.

1. INTRODUCTION

Let F_n be the free group of finite rank n . For any endomorphism α of F_n the *fixed point subgroup* of α is

$$\text{Fix}(\alpha) = \{x \in F_n \mid \alpha(x) = x\}.$$

More generally, if S is a set of endomorphisms of F_n , the *fixed point subgroup* of S is $\text{Fix}(S) = \bigcap_{\alpha \in S} \text{Fix}(\alpha)$.

Dyer and Scott showed in [19] that if α is an automorphism of finite order of F_n , then $\text{Fix}(\alpha)$ is a free factor of F_n , in particular the rank of $\text{Fix}(\alpha)$ does not exceed n . Scott later (1978) conjectured that

$$\text{rk Fix}(\alpha) \leq n$$

for any automorphism of F_n .

Gersten [24, 25], Goldstein and Turner [27], and Cooper [17] independently proved a weaker form of this conjecture that the group $\text{Fix}(\alpha)$ is finitely generated. In [36], Thomas generalized this result for an arbitrary set of automorphisms of F_n . In [35], Stallings proved that the equalizer subgroup

$$\text{Eq}(\varphi, \psi) := \{x \in F_n \mid \varphi(x) = \psi(x)\}$$

is finitely generated for any two monomorphisms φ, ψ from F_n into a free group G . Goldstein and Turner [26] showed that the same holds in case where φ is a homomorphism and ψ is a monomorphism from F_n to F_n .

In the seminal paper [7], Bestvina and Handel proved the Scott conjecture for automorphisms of free groups completely. The relative train track technique developed in this paper enabled the solution of a sequel of difficult problems about automorphisms of free groups, see [4, 5, 6, 15, 11, 20, 13, 12]. Later Gaboriau, Levitt and Lustig [23] gave a dendrological proof of the Scott conjecture. A stronger inequality, which also takes into account infinite fixed words, is given in [21]. Another dendrological proof was given by Sela in [33]. A short dendrological proof of the Scott conjecture based on papers [23], [33], and [32] is given in the survey [3] of Bestvina on $\mathbb{R}$ -trees.

Imrich and Turner [28] proved that $\text{rk Fix}(\varphi) \leq n$ for an arbitrary endomorphism φ of F_n . Dicks and Ventura [18] proved that, $\text{rk Fix}(S) \leq n$ for any set S of injective endomorphisms of F_n . Bergman, using derivations in group rings, proved that the same holds for an arbitrary set of endomorphisms of F_n .

However, the problem of finding a basis of $\text{Fix}(\alpha)$ has been open for almost 20 years. It has been solved in three special cases: for positive automorphisms in the paper [16] of Cohen and Lustig, for special irreducible automorphisms in the paper of Turner [37, Proposition B], and for all automorphisms of F_2 in the paper of Bogopolski [9].

In 1999, Maslakova, a former PhD student of the first named author, attempted to solve this problem in general case. However, her proof published in [30] was not complete. An improved, but still incomplete proof was given in her PhD thesis [31]. So, we have decided to fix all mistakes and give a correct proof. The main result of this paper is the following.

Theorem 1.1. *For any automorphism α of a free group F of finite rank, one can compute a basis of the fixed point subgroup $\text{Fix}(\alpha)$.*

As in [7], we use the relative train track techniques for automorphisms of free groups. A relative train track for α is a homotopy equivalence $f : \Gamma \rightarrow \Gamma$ of a finite graph Γ with certain good properties (see Section 4). We use f to define an auxiliary graph D_f (first introduced in [26]); the fundamental group of one of the components of D_f , denoted $D_f(\mathbf{1}_v)$, can be identified with $\text{Fix}(\alpha)$ (see Section 5). Thus, to compute a basis of $\text{Fix}(\alpha)$, we need to construct the core $\text{Core}(D_f(\mathbf{1}_v))$ of this component.

At all but finite number of the vertices of D_f there is a preferable outgoing direction. This determines a flow on almost all of D_f . The inverse automorphism α^{-1} determines its own flow on almost all of D_f which does not coincide with the flow for α outside a finite subgraph. According to [37] (see also [16]), there is a procedure for constructing a part of $\text{Core}(D_f)$ which contains $\text{Core}(D_f(\mathbf{1}_v))$ if the latter is non-contractible: one should start from a finite number of computable exceptional edges and follow the first flow for sufficiently long. Theoretically we could arrive at a dead vertex, or get a loop, or arrive at a vertex where two rays of this flow meet, or none of these may occur. To convert this procedure into an algorithm, we must detect at the beginning, which possibility occurs. Also we must solve a membership problem for vertices and certain subgraphs of D_f . We overcome these difficulties in this paper.

2. PRELIMINARIES

Let Γ be a finite connected graph, Γ^0 be the set of its vertices, Γ^1 be the set of its edges. The inverse edge to E is denoted by $\overline{E}$. Sometimes we will consider the geometric realization of Γ where each edge is isometric to the real segment $[0, 1]$. For simplicity, we will denote this realization also by Γ .

A *path* in Γ is a continuous map $\tau : [0, 1] \rightarrow \Gamma$ which is locally injective on $\tau^{-1}(\Gamma \setminus \Gamma^0)$. Let $\alpha(\tau)$ and $\omega(\tau)$ be the initial and the terminal points of τ . These points are not necessarily vertices of Γ . The inverse path to τ is denoted by $\bar{\tau}$ and the length of τ by $l(\tau)$. A path whose image is a point u is called *trivial* and is denoted by $\mathbf{1}_u$. A *combinatorial representative* of a nontrivial path is a sequence $E_1 E_2 \dots E_n$ of (segments of) edges which it passes; if $n = 1$, then E_1 is a subsegment of an edge; if $n \geq 2$, then E_i are edges for $1 < i < n$ and E_1, E_n are some terminal and initial segments of some edges. Sometimes we will identify paths with their combinatorial representatives.

An *edge-path* in Γ is a path of the form $E_1 E_2 \dots E_n$ where E_i are edges of Γ , or a path of the form $\mathbf{1}_u$ where u is a vertex of Γ . In the latter case the edge-path is called *trivial*. Let $P(\Gamma)$ be the set of all nontrivial edge-paths in Γ .

A path τ is called *reduced* if it does not contain nontrivial subpaths of the form $\mu\bar{\mu}$. By $[\tau]$ we denote the reduced path in Γ which is homotopic to τ relative to its endpoints. By $[[\tau]]$ we denote the class of paths in Γ which are homotopic to τ relative to the endpoints of τ . For two paths τ, μ , we write $\tau = \mu$ if these paths are homotopic and $\tau \equiv \mu$ if they coincide.

For two paths τ, μ with $\omega(\tau) = \alpha(\mu)$, we denote their concatenation by $\tau\mu$ or $\tau \cdot \mu$. A vertex or an edge of Γ can occur several times in a given path; it will be clear from a context which occurrence do we mean.

Let $\mathcal{HE}$ be the set of all homotopy equivalences $f : \Gamma \rightarrow \Gamma$ such that Γ is a finite connected graph, $f(\Gamma^0) \subseteq \Gamma^0$ and the map $f|_{\Gamma \setminus \Gamma^0}$ is locally injective.

Let $f : \Gamma \rightarrow \Gamma$ be a homotopy equivalence from $\mathcal{HE}$. Then for every path τ in Γ the map $f \circ \tau$ is also a path in Γ . We denote this path by $f(\tau)$. In particular, if we consider an edge $E \in \Gamma^1$ as a path, then $f(E)$ is a nontrivial reduced edge-path.

The *combinatorial representative* of f is the map $\Gamma^1 \rightarrow P(\Gamma)$ which sends each edge $E \in \Gamma^1$ to the combinatorial representative of the path $f(E)$. Observe that the set of combinatorial representatives of homotopy equivalences with the above properties is enumerable.

The *norm* of f is the number

$$\|f\| := \max\{l(f(E)) \mid E \text{ is an edge of } \Gamma\}.$$

If v is a distinguished vertex of Γ , we write $f_* : \pi_1(\Gamma, v) \rightarrow \pi_1(\Gamma, f(v))$ for the isomorphism given by $[[x]] \mapsto [[f(x)]]$, $x \in \pi_1(\Gamma, v)$. Observe that if $f : \Gamma \rightarrow \Gamma$ and $g : \Gamma \rightarrow \Gamma$ are two homotopy equivalences from $\mathcal{HE}$ with the same combinatorial

representatives, then $f_* = g_*$. Sometimes we will identify homotopy equivalences with their combinatorial representatives.

We use the following rule for composition of two maps: $x(\varphi\psi) = (x\varphi)\psi$.

3. OUTLINE OF THE PROOF

In this section we give a sketch of the proof of the main theorem. Some of definitions we use here are given in the following sections. At this point, it suffices to know the definition of a relative train track (see Section 4).

A. Let α be an automorphism of a free group F of finite rank.

Definition 3.1. We say that α is *realized* by a homotopy equivalence $f : \Gamma \rightarrow \Gamma$, where Γ is a finite connected graph, if there is a vertex v in Γ fixed by f and an isomorphism $j : F \rightarrow \pi_1(\Gamma, v)$ such that the automorphism $j^{-1}\alpha j$ of the group $\pi_1(\Gamma, v)$ coincides with the induced automorphism f_* . We will also say that f *represents* α .

Observe that for computing a basis of $\text{Fix}(\alpha)$, it suffices to compute a basis of $\text{Fix}(\alpha^n)$ for some $n \geq 1$ (see Corollary 4.11). So we can replace α by an appropriate power of α if needed. Using this and Theorem 4.6, we may assume that α is algorithmically realized by a relative train track $f : (\Gamma, v) \rightarrow (\Gamma, v)$ with the additional property (Pol). Let $\emptyset = G_0 \subset \dots \subset G_N = \Gamma$ be the fixed maximal filtration for f ; this gives us exponential, polynomial, or zero strata $H_i := \text{cl}(G_i \setminus G_{i-1})$, $i = 1, \dots, N$. In Section 4 we recall that for each exponential stratum H_r , there exists a real number $\lambda_r > 1$ and a pseudo-metric L_r on G_r with the following properties:

- if E is an edge in G_r , then $L_r(E) > 0$ if and only if E is an edge in H_r ;
- if p is a path in G_r , then $L_r(f(p)) = \lambda_r L_r(p)$;
- if p is a reduced r -legal path in G_r , then $L_r([f(p)]) = \lambda_r L_r(p)$.

B. Here we describe some technical notions from Section 7 including the notion of an r -cancellation area. Using this notion, we improve f in Section 9.

To understand this part, we advise first to read the definitions of a legal and an illegal r -turns in Section 4.

Let τ be a reduced edge-path in Γ and y be an occurrence of a vertex in τ . Let p_0, q_0 be paths in Γ such that $\tau \equiv \bar{p}_0 q_0$ and $y = \alpha(p_0) = \alpha(q_0)$. Then $[f(\tau)]$ is the result of reducing of the product $[f(\bar{p}_0)] \cdot [f(q_0)]$. Let $[f(\bar{p}_0)] \equiv \bar{p}_1 \bar{r}$ and $[f(q_0)] \equiv r q_1$ for some paths p_1, q_1, r such that $[f(\tau)] \equiv \bar{p}_1 q_1$. Then the occurrence $y_1 := \alpha(\bar{p}_1) = \alpha(q_1)$ in $[f(\tau)]$ is called the *1-successor* of y . Inductively, one can define y_k , the *k-successor* of y for $k \geq 1$.

For the rest of this subsection, we suppose additionally that τ lies in G_r , where H_r is an exponential stratum. The occurrence y in τ is called an *r-cancellation*

point in τ if the pair (A, B) consisting of the first edges of p_0 and q_0 is an illegal r -turn. One can easily check, whether y is an r -cancellation point in τ .

In Section 7, we consider the case where y is a unique r -cancellation point in τ . In this case we say that y is *non-deletable* if its k -successor y_k is an r -cancellation point in $[f^k(\tau)]$ for each $k \geq 1$. By Proposition 7.9, one can check whether y is non-deletable in τ .

Now we suppose that y is a unique r -cancellation point in τ and that y is non-deletable. In Definition 7.13 we define the *r -cancellation area* of y in τ . Roughly speaking this is a subpath $A(\tau, y)$ of τ which contains y and consists of all points of τ which cancel out by reducing the product $[f^k(\bar{p}_0)] \cdot [f^k(q_0)]$ for appropriate $k \geq 0$. We stress that $A(\tau, y)$ is not necessarily an edge-path. If τ is fixed, we shorten the notation to $A(y)$.

The r -cancellation areas possess the following properties:

- 1) The endpoints of each r -cancellation area $A(\tau, y)$ lie in r -edges of Γ (i.e. in edges which belong to H_r).
- 2) One can compute the number $L_r(A(\tau, y))$.
- 3) One can compute a number $R = R(f)$ such that each r -cancellation area is divided by the corresponding non-deletable r -cancellation point into two subpaths which are contained in edge-paths of l -length at most R .
- 4) There exists only finitely many r -cancellation areas. These areas can be computed.
- 5) If A is an r -cancellation area in τ , then $[f(A)]$ is an r -cancellation area in $[f(\tau)]$.

In Section 9, we use a subdivision to obtain a new relative train track $f' : \Gamma' \rightarrow \Gamma'$ representing α and satisfying the following additional condition.

(RTT-iv) There is a computable natural number $P = P(f)$ such that for each exponential strata H_r and each r -cancellation area A of f , the r -cancellation area $[f^P(A)]$ is an edge-path.

We describe briefly how we do that. Let H_r be an exponential strata for the relative train track $f : \Gamma \rightarrow \Gamma$ and let $A_1, \dots, A_s$ be all r -cancellation areas of f . An endpoint u of A_i is called an *r -subdivision point* if u is not a vertex (hence u lies in the interior of an edge from H_r) and $f^n(u)$ is not a vertex for all natural n .

By Property 4), we can construct the graph Γ' obtained from Γ by subdivision at all r -subdivision points for all exponential strata H_r . Property 5) implies that the set of these endpoints is f -invariant. Let $f' : \Gamma' \rightarrow \Gamma'$ be the map induced by $f : \Gamma \rightarrow \Gamma$. We also define a natural filtration $\emptyset = G'_0 \subset \dots \subset G'_N = \Gamma'$ by saying that G'_i consists of all edges resulting from G_i by the subdivision. It is possible that this filtration is not maximal. A strata H'_i is called exponential, polynomial, or zero if the strata H_i is exponential, polynomial, or zero, respectively. For each

exponential stratum H'_r , let L'_r be the pseudo-metric on G'_r induced by L_r . It is possible that $L'_r(e) = 0$ for some edge $e \in H'_r$. However, this will not destroy the further proof.

One can show that the homotopy equivalence $f' : \Gamma' \rightarrow \Gamma'$ satisfies the conditions (RTT-i) – (RTT-iii) in the definition of a relative train track. Therefore f' is called the *subdivided relative train track* associated with f . The advantage of f' is that it additionally satisfies the condition (RTT-iv).

Moreover, the map f' also represents the automorphism α and satisfies the property (Pol). We will work with f' starting from Section 10.

C. The following definition generalizes the above definition of a single non-deletable r -cancellation point in a path τ . Let τ be a reduced edge-path in G_r and let $y_1, \dots, y_k$ be all r -cancellation points in τ . We say that these points are *non-deletable* in τ if the number of r -cancellation points in $[f^i(\tau)]$ is equal to k for each $i \geq 0$.

By Propositions 8.4 and 7.9, we can check, whether the points $y_1, \dots, y_k$ are non-deletable in τ . If they are non-deletable, the path τ is called *stable*. By Proposition 8.2, one can algorithmically find a number $s \geq 0$ such that the path $[f^s(\tau)]$ is stable.

If τ is stable and $y_1, \dots, y_k$ are all non-deletable r -cancellation points in τ , then we can write τ in the form $\tau \equiv b_0 A_1 b_1 \dots A_k b_k$, where the paths b_j are r -legal or trivial and $A_1, \dots, A_k$ are r -cancellation areas corresponding to $y_1, \dots, y_k$ in τ . Such expression is called the *A-decomposition* of τ .

By Remark 8.7, if τ has the A -decomposition $\tau \equiv b_0 A_1 b_1 \dots A_k b_k$, then for every $i \geq 1$, the path $[f^i(\tau)]$ has the A -decomposition $[f^i(\tau)] \equiv b_0^i \cdot A_1^i \cdot b_1^i \cdot \dots \cdot A_k^i \cdot b_k^i$, where $b_j^i \equiv [f^i(b_j)]$ and $A_j^i \equiv [f^i(A_j)]$ for all possible j .

Consider τ as a path in the subdivided relative train track Γ' and let $P = P(f')$ be the constant from (RTT-iv). Then the A -decomposition of $[f^P(\tau)]$ has the property that all the subpaths A_j^P and b_j^P are edge-paths in Γ' . This observation explains why we prefer to work with the subdivided relative train track f' rather than with f .

D. Let Γ be a finite connected graph with a distinguished vertex v and let $f : \Gamma \rightarrow \Gamma$ be a homotopy equivalence such that $f(\Gamma^0) \subseteq \Gamma^0$, f is locally injective on $\Gamma \setminus \Gamma^0$, and $f(v) = v$. Thus, f is not necessarily a (subdivided) relative train track. In Section 5 and here we define a graph D_f and describe a procedure which helps to compute a basis of the group

$$\overline{\text{Fix}}(f) := \{[[p]] \in \pi_1(\Gamma, v) \mid f(p) = p\}.$$

In Sections 9-15, we will convert this procedure into an algorithm in case, where f is a subdivided relative train track satisfying the property (Pol). From this we will deduce an algorithm for computing $\text{Fix}(\alpha)$.

A definition of the graph D_f . An f -path in Γ is an edge-path μ in Γ (possibly trivial) such that f maps the initial point of μ to the terminal point of μ . Thus, if μ is an f -path in Γ , the path $\mu f(\mu)$ is well defined. The vertices of the graph D_f are reduced f -paths. A vertex μ of D_f is called *trivial* if $\mu = \mathbf{1}_u$ for some vertex u of Γ fixed by f . Two vertices μ and τ in D_f are connected by an edge (from μ to τ) with label E if E is an edge in Γ outgoing from the same vertex as μ and we have $[\overline{E}\mu f(E)] = \tau$.

We set $\widehat{f}(\mu) := [\overline{E}\mu f(E)]$ if E is the first edge of the f -path μ (in particular, μ must be nontrivial). Clearly, μ and $\widehat{f}(\mu)$ are connected by an edge with the label E in D_f . The direction of this edge is called *preferable* at μ . Preferable directions at all nontrivial vertices determine a flow in D_f . Starting at μ and moving along this flow, we get vertices $\mu = \mu_1, \mu_2, \dots$, where $\mu_{i+1} = \widehat{f}(\mu_i)$, $i \geq 1$. These vertices together with the directed edges we pass form a subgraph in D_f which we call the μ -subgraph. The μ -subgraph is either a finite segment, or a finite segment with a cycle, or a ray (see Figure 5).

An edge e connecting two vertices u, w in D_f is called *repelling* if the direction of e is not preferable at u and the direction of the opposite edge $\bar{e}$ is not preferable at w . Endpoints of repelling edges are called *repelling vertices*. By Proposition 5.2, there exist only finitely many repelling edges in D_f and they can be algorithmically found.

E. Recall that v is the distinguished vertex of Γ and $f(v) = v$. Since $\mathbf{1}_v$ is an f -path, we can consider $\mathbf{1}_v$ as a vertex of D_f . Let $D_f(\mathbf{1}_v)$ be the component of D_f containing $\mathbf{1}_v$. The fundamental group $\pi_1(D_f(\mathbf{1}_v), \mathbf{1}_v)$ can be identified with $\overline{\text{Fix}}(f)$, see Lemma 5.3.

A component of D_f is called *repelling* if it contains at least one repelling edge. Let $C_1, \dots, C_n$ be all repelling components of D_f . For each C_i , let $\text{CoRe}(C_i)$ be a core of C_i which contains all repelling edges of C_i . We set $C_f := \cup_{i=1}^n C_i$ and $\text{CoRe}(C_f) := \cup_{i=1}^n \text{CoRe}(C_i)$.

By Proposition 5.6, to compute a basis of $\pi_1(D_f(\mathbf{1}_v), \mathbf{1}_v)$, it suffices to construct $\text{CoRe}(C_f)$ and to decide, whether the vertex $\mathbf{1}_v$ lies in the μ -subgraph for some repelling vertex μ .




Figure 1.

An example of a graph D_f with three repelling components. One of them is contractible.

F. It turns out that $CoRe(C_f)$ is contained in the union of the repelling edges and the μ -subgraphs, where μ runs over the set of repelling vertices. So, to construct $CoRe(C_f)$, it suffices to do the following:

- (1) Compute repelling edges.
- (2) For each repelling vertex μ determine, whether the μ -subgraph is finite or not.
- (3) Compute all elements of all finite μ -subgraphs from (2).
- (4) For each two repelling vertices μ and τ with infinite μ - and τ -subgraphs determine, whether these subgraphs intersect.
- (5) If the μ -subgraph and the τ -subgraph from (4) intersect, find their first intersection point and compute their initial segments up to this point.

As it was mentioned above, step (1) can be done algorithmically. Using Propositions 5.14 and 5.15, we show in Section 5 that steps (2)-(5) can be done algorithmically if the following two problems are solvable:

Problem 1. Given two vertices μ and τ of the graph D_f , verify whether τ is contained in the μ -subgraph.

Problem 2. Given a vertex μ of the graph D_f , determine whether the μ -subgraph is finite or not.

In the part G, we sketch a solution of these problems in the case where f is a subdivided relative train track satisfying the property (Pol). Hence, for such f we can compute a basis of $\overline{\text{Fix}}(f)$. By the part B, we may assume that α is realized by such f . Using this, we can compute a basis of $\text{Fix}(\alpha)$.

G. From now on and to the end of this section, we assume that $f : (\Gamma, v) \rightarrow (\Gamma, v)$ is a subdivided relative train track satisfying the property (Pol). Let $\emptyset = G_0 \subset \dots \subset G_N = \Gamma$ be the filtration associated with f .

To solve Problems 1 and 2, we investigate μ -subgraphs in D_f in details. The difficulty is that we cannot control cancelations in passing from $\mu \equiv E_1 E_2 \dots E_s$ to $\widehat{f}(\mu) \equiv [E_2 \dots E_s \cdot f(E_1)]$ and, more generally, from $\widehat{f}^i(\mu)$ to $\widehat{f}^{i+1}(\mu)$, $i \geq 1$. However, we will show that *if the μ -subgraph is infinite, it contains a vertex v_0 which is in some sense perfect* (see Definitions 10.1, 10.3, and 14.1 where we define r -perfect, A -perfect, and E -perfect f -paths, respectively; see also Propositions 13.4 and 14.2 for the proof).

If v_0 is r -perfect or A -perfect, there is no cancelation in passing from v_0 to $\widehat{f}(v_0)$. Some other good properties of perfect vertices are collected in Propositions 10.2, 10.4, and 14.2.

Observe that

- if the μ -subgraph is infinite, it contains infinitely many perfect vertices;
- for any vertex u of D_f , one can check, whether u is perfect and if yes, one can detect the kind of perfectness of u .

Hence, moving along the μ -subgraph, we either detect that the μ -subgraph is finite, or we find a perfect vertex v_0 in the μ -subgraph of sufficiently large l -length (we need v_0 with $l(v_0) \geq K$, where $K = K(f)$ is a computable constant from Proposition 9.5. 8)). Note that in the second case we still have to decide whether the μ -subgraph is infinite. In G1 – G3, we briefly describe how we do this depending on the kind of perfectness of v_0 . The full proof is given in Section 15.

G1. If v_0 is an r -perfect path in G_r , then the stratum H_r is exponential. Let $\lambda_r > 1$ be the Perron-Frobenius eigenvalue corresponding to H_r . Then, by Lemma 15.1 we have the following:

- (1) $L_r(\widehat{f}^{i+1}(v_0)) \geq L_r(\widehat{f}^i(v_0)) > 0$ for all $i \geq 0$.
- (2) There exist computable natural numbers $m_1 < m_2 < \dots$, such that $L_r(\widehat{f}^{m_i}(v_0)) = \lambda_r^i L_r(v_0)$ for all $i \geq 1$.

Hence, if v_0 is r -perfect, then the v_0 -subgraph is infinite, and so the μ -subgraph is infinite, which solves Problem 2. Also Problem 1 can be easily solved with the help of L_r .

G2. If v_0 is an A -perfect path in G_r and r is minimal, then the stratum H_r is exponential as well. By Proposition 10.4, we can algorithmically find a finite number of A -perfect f -paths $\mu_{0,1}, \dots, \mu_{0,k}$, where $v_0 = \mu_{0,1}$, which, considered as vertices of D_f , lie in the μ -subgraph and satisfy the following properties:

- (1) For any $1 \leq j \leq k$ and $i \geq 0$, the path $\mu_{i,j} := [f^i(\mu_{0,j})]$ is an A -perfect f -path.
- (2) For any $1 \leq j \leq k$ and $i \geq 0$, the vertex $\mu_{i,j}$ of D_f lies in the $\mu_{0,1}$ -subgraph. Moreover $\mu_{i,j} \equiv \widehat{f}^{m_{i,j}}(\mu_{0,1})$ for some $m_{i,j}$ satisfying $m_{i,j} < m_{i,j+1}$ and $m_{i,k} < m_{i+1,1}$.
- (3) If σ is a vertex of the $\mu_{0,1}$ -subgraph such that the f -path σ is A -perfect, then σ coincides with $\mu_{i,j}$ for some i, j .

This implies that we can solve Problem 2 if we answer the following question: Does there exist $0 \leq p < q$ such that $[f^p(\mu_{0,1})] = [f^q(\mu_{0,1})]$? This question can be answered with the help of Corollary 4.9.

We will reduce Problem 1 to the case where τ is A -perfect. In this case we will reduce Problem 1 to the following question: Does there exist $p \geq 0$ such that $[f^p(\mu_{0,j})] = \tau$ for some $1 \leq j \leq k$? This questions can be answered with the help of Corollary 4.8.

Both corollaries follow from Theorem 4.7 which was proved by P. Brinkmann (see [13, Theorem 0.1]).

G3. If v_0 is an E -perfect f -path in G_r and r is minimal, then H_r is a polynomial stratum. Let E be the unique (up to inversion) edge of H_r . We write the f -path v_0 in the form $v_0 \equiv E_1 b_1 E_2 \dots E_k b_k$, where $E_1, \dots, E_k \in \{E, \overline{E}\}$ and $b_1, \dots, b_k$ are paths in G_{r-1} .

By Proposition 14.2, we can algorithmically find a finite number of E -perfect f -paths $\mu_{0,1}, \dots, \mu_{0,k}$, where $v_0 = \mu_{0,1}$, which, considered as vertices of D_f , lie in the μ -subgraph and satisfy the properties analogous to (1)-(3) from Case G2. The only difference is that the A -perfectness must be replaced by the E -perfectness and the definition of $\mu_{i,j}$ must be replaced by the following: $\mu_{i,j} := [\overline{c_{i,E_j}} f^i(\mu_{0,j}) f(c_{i,E_j})]$, where c_{i,E_j} are paths defined at the beginning of Section 14. This and some rewriting trick enable us to solve Problems 1 and 2 with the help of Corollaries 4.8 and 4.9 similarly to Case G2 (see Sections 14 and 15).

4. A RELATIVE TRAIN TRACK FOR AN AUTOMORPHISM OF A FREE GROUP OF FINITE RANK

First we recall the definition of a relative train track from [7]. Let $f : \Gamma \rightarrow \Gamma$ be a homotopy equivalence such that $f(\Gamma^0) \subseteq \Gamma^0$ and the map $f|_{\Gamma \setminus \Gamma^0}$ is locally injective. As it was mentioned in Section 2, for every edge $E \in \Gamma^1$, we can consider $f(E)$ as a nontrivial reduced edge-path.

A *turn* in Γ is an unordered pair of edges of Γ originating at a common vertex. A turn is *nondegenerate* if these edges are distinct, and it is *degenerate* otherwise. The map $f : \Gamma \rightarrow \Gamma$ induces a map $Df : \Gamma^1 \rightarrow \Gamma^1$ which sends each edge $E \in \Gamma^1$ to the first edge of the path $f(E)$. This induces a map Tf on turns in Γ by the rule $Tf(E_1, E_2) = (Df(E_1), Df(E_2))$. A turn (E_1, E_2) is *legal* if the turns $(Tf)^n(E_1, E_2)$ are nondegenerate for all $n \geq 0$; a turn is *illegal* if it is not legal. An edge-path $E_1 E_2 \dots E_m$ in Γ is *legal* if all its turns $(\overline{E}_i, E_{i+1})$ are legal. Clearly, a legal edge-path is reduced.

From each pair of mutually inverse edges of the graph Γ we choose one edge. Let $\{E_1, \dots, E_k\}$ be the ordered set of chosen edges. The *transition matrix* of the map f (with respect to this ordering) is the matrix $M(f)$ of the size $k \times k$ such that the ij^{th} entry of $M(f)$ is equal to the total number of occurrences of the edges E_i and $\overline{E}_i$ in the path $f(E_j)$.

A *filtration* for $f : \Gamma \rightarrow \Gamma$ is an increasing sequence of (not necessarily connected) f -invariant subgraphs $\emptyset = G_0 \subset \dots \subset G_N = \Gamma$. The subgraph $H_i = \text{cl}(G_i \setminus G_{i-1})$ is called the *i -th stratum*. Edges in H_i are called *i -edges*. A turn with both edges in H_i is called an *i -turn*. A turn with one edge in H_i and another in G_{i-1} is called *mixed* in (G_i, G_{i-1}) . We assume that the edges of Γ are ordered so that the edges from H_i precede the edges from H_{i+1} . The edges from H_i define a square submatrix M_i of $M(f)$.

If the filtration is maximal, then each matrix M_i is irreducible. If M_i is nonzero and irreducible, then it has the associated Perron-Frobenius eigenvalue $\lambda_i \geq 1$. If $\lambda_i > 1$, then the stratum H_i is called *exponential*. If $\lambda_i = 1$, then the stratum H_i is called *polynomial*. In this case M_i is a permutation matrix, hence for every edge $E \in H_i$ the path $f(E)$ contains exactly one edge of H_i , all other edges of $f(E)$ lie in G_{i-1} . A stratum H_i is called a *zero stratum* if M_i is a zero matrix. In this case $f(E)$ lies in G_{i-1} for every edge $E \in H_i$.

Let $f : \Gamma \rightarrow \Gamma$ be a homotopy equivalence such that $f(\Gamma^0) \subseteq \Gamma^0$ and the map $f|_{\Gamma \setminus \Gamma^0}$ is locally injective. The map f is called a *relative train track* if there exists a maximal filtration for f such that each exponential stratum H_r of this filtration satisfies the following conditions:

- (RTT-i) Df maps the set of edges of H_r to itself; in particular all mixed turns in (G_r, G_{r-1}) are legal.
- (RTT-ii) If $\rho \subset G_{r-1}$ is a reduced nontrivial edge-path with endpoints in $H_r \cap G_{r-1}$, then $[f(\rho)]$ is a nontrivial edge-path with endpoints in $H_r \cap G_{r-1}$.
- (RTT-iii) For each legal edge-path $\rho \subset H_r$, the path $f(\rho)$ does not contain any illegal turns in H_r .

A nontrivial reduced edge-path in G_r is called *r-legal* if it has no illegal turns in H_r . A nontrivial reduced path ρ in G_r is called *r-legal* if the minimal edge-path containing ρ is *r-legal*.

Proposition 4.1. (see [7, Lemma 5.8]) *Suppose that $f : \Gamma \rightarrow \Gamma$ is a relative train track and H_r is an exponential stratum of Γ . Let ρ be a reduced *r-legal* path:*

$$\rho \equiv b_0 \cdot a_1 \cdot b_1 \cdot \dots \cdot a_k \cdot b_k,$$

where $k \geq 1$, $a_1, \dots, a_k$ are paths in H_r , and $b_0, \dots, b_k$ are paths in G_{r-1} , and all these paths except maybe b_0 and b_k are nontrivial. Then

$$[f(\rho)] \equiv [f(b_0)] \cdot f(a_1) \cdot [f(b_1)] \cdot \dots \cdot f(a_k) \cdot [f(b_k)]$$

and this path is *r-legal*. Moreover, for all $i \geq 1$ we have

$$[f^i(\rho)] \equiv [f^i(b_0)] \cdot [f^i(a_1)] \cdot [f^i(b_1)] \cdot \dots \cdot [f^i(a_k)] \cdot [f^i(b_k)]$$

and these paths are *r-legal*.

The *r*-length function L_r . Let H_r be an exponential stratum and let $\vec{v}$ be a positive vector satisfying $\vec{v}M_r = \lambda_r \vec{v}$. Since M_r is an integer matrix, we can choose $\vec{v}$ so that the coordinates of $\vec{v}$ are rational functions of λ_r over $\mathbb{Q}$. If E_i is the i^{th} edge of H_r , define $L_r(E_i) = v_i$; if E is an edge of G_{r-1} , define $L_r(E) = 0$. For an arbitrary edge-path τ in G_r , we define its *r-length* $L_r(\tau)$ as the sum of *r*-lengths of edges of τ . Then we have $L_r(f(E)) = \lambda_r L_r(E)$.

Lemma 5.10 in [7] claims that L_r can be extended to a function $L_r(\rho)$ for paths $\rho \subset G_r$ so that we will have $L_r(f(\rho)) = \lambda_r L_r(\rho)$ for any *r-legal* path ρ . (Note that the proof of this lemma shows that this formula holds for any path $\rho \subset G_r$.) This and Proposition 4.1 imply that for every exponential stratum H_r and every reduced *r-legal* path $\rho \subset G_r$ we have $L_r([f(\rho)]) = \lambda_r L_r(\rho)$.

The function L_r has else one useful property: if ρ is a nontrivial initial or terminal segment of an edge in H_r , then $L_r(\rho) > 0$.

In [7, Theorem 5.12]), Bestvina and Handel proved that for every outer automorphism $\mathcal{O}$ of F_n , there exists a *stable* relative train track $f : \Gamma \rightarrow \Gamma$ representing $\mathcal{O}$. Their proof does not give an algorithm to construct such train track. However, one can extract from the proof an algorithm constructing a relative train track for $\mathcal{O}$. We formulate this more precisely for automorphisms of free groups.

Theorem 4.2. (see [7, Theorem 5.12]) *Let F be a free group of finite rank. For each automorphism α of F , one can algorithmically construct a finite connected graph Γ and a relative train track $f : \Gamma \rightarrow \Gamma$, and indicate a vertex $v \in \Gamma^0$, a path $p \subset \Gamma$ from v to $f(v)$, and an isomorphism $i : F \rightarrow \pi_1(\Gamma, v)$ such that the automorphism $i^{-1}\alpha i$ of the group $\pi_1(\Gamma, v)$ coincides with the map given by the rule*

$$[[x]] \mapsto [[p]]f_*([[x]])[[\bar{p}]],$$

where $[[x]] \in \pi_1(\Gamma, v)$.

Remark. There is also a brute force way to construct a relative train track (abbreviated RRT) representing α in the sense of the above theorem. The idea is to enumerate all combinatorial RRT and then find one which represents α .

We explain why the set of all combinatorial relative train tracks is enumerable. First, one can enumerate the combinatorial representatives of all homotopy equivalences $f : \Gamma \rightarrow \Gamma$ with the property that Γ is a finite connected graph, $f(\Gamma^0) \subseteq \Gamma^0$, and the map $f|_{\Gamma \setminus \Gamma^0}$ is locally injective. For any given f , one can enumerate all possible maximal filtrations. Then one can check, given f with a maximal filtration, whether they satisfy (RTT-i) – (RTT-iii). The verification of (RTT-i) is trivial, the verification of (RTT-iii) can be reduced to that for edges of H_r and so is also trivial. Thus, we may assume that (RTT-i) and (RTT-iii) are satisfied. In particular, the set of vertices $H_r \cap G_{r-1}$ is f -invariant. Now we explain how to verify (RTT-ii).

Given a pair of vertices u, u' in $H_r \cap G_{r-1}$, we have to verify, whether (RTT-ii) is fulfilled for $\rho \subset G_{r-1}$ connecting u and u' . Since f is a homotopy equivalence, (RTT-ii) is fulfilled in case $u = u'$. So, we may assume that $u \neq u'$.

If u, u' lie in different components of G_{r-1} , there is no such ρ and (RTT-ii) is trivially fulfilled. So, we assume that there is a path in G_{r-1} from u to u' . Choose one of them and denote it by $\tau_{u,u'}$.

If $f(u) \neq f(u')$, then every edge-path $\rho \subset G_{r-1}$ with endpoints u, u' satisfies (RTT-ii). Suppose that $f(u) = f(u')$. Since $f(\tau_{u,u'})$ is a loop, we can find a reduced loop $\tau_u \subset \Gamma$ based at u such that $[f(\tau_{u,u'})] = [f(\tau_u)]$. We claim that (RTT-ii) is fulfilled for any $\rho \subset G_{r-1}$ connecting u and u' if and only if $\tau_u \not\subset G_{r-1}$. Indeed, suppose that $[f(\rho)]$ is trivial. Then $[f(\tau_{u,u'}\bar{\rho})] = [f(\tau_{u,u'})] = [f(\tau_u)]$, hence the loops $[\tau_{u,u'}\bar{\rho}]$ and τ_u coincide, in particular $\tau_u \subset G_{r-1}$. Conversely, if $\tau_u \subset G_{r-1}$, then $\rho := [\bar{\tau}_u\tau_{u,u'}]$ lies in G_{r-1} and nontrivial, but the path $[f(\rho)]$ is trivial.

Thus, we can enumerate all combinatorial relative train tracks. For a given combinatorial relative train track $f : \Gamma \rightarrow \Gamma$, we can enumerate all vertices $v \in \Gamma$ and paths p from v to $f(v)$, and isomorphisms $i : F \rightarrow \pi_1(\Gamma, v)$. For every such f, v, p, i , we can verify, whether the formula in Theorem 4.2 determines the isomorphism $i^{-1}\alpha i$. Proceeding diagonally, we can find the desired relative train track. $\square$

Below we will replace the map $f : \Gamma \rightarrow \Gamma$ by an iterate. Doing that, we must be careful: a positive power of f can map an edge of Γ to a non-reduced edge-path. Thus, we deem the phrase *passing to a positive power of f* , to mean that for fixed $k \in \mathbb{N}$, we pass from consideration of $f : \Gamma \rightarrow \Gamma$ to consideration of the map $f^k_\# : \Gamma \rightarrow \Gamma$ that sends each edge E in Γ to the reduced edge-path $[f^k(E)]$.

Passing to an appropriate positive power of f , refining the maximal filtration, and using subdivisions of edges of polynomial strata, one can prove the following variation of Theorem 4.2 (which is a part of Theorem 5.1.5 in [4]).

Theorem 4.3. *Let F be a free group of finite rank. For any automorphism α of F , one can algorithmically compute a natural number n , construct a finite connected graph Γ and a relative train track $f : \Gamma \rightarrow \Gamma$, and indicate a vertex $v \in \Gamma^0$, a path $p \subset \Gamma$ from v to $f(v)$, and an isomorphism $i : F \rightarrow \pi_1(\Gamma, v)$ such that the automorphism $i^{-1}\alpha^n i$ of the group $\pi_1(\Gamma, v)$ coincides with the map given by the rule*

$$[[x]] \mapsto [[p]]f_*([x])[[\bar{p}]],$$

where $[[x]] \in \pi_1(\Gamma, v)$, and the following condition is satisfied:

(Pol) *Every polynomial stratum H_r consists of only two mutually inverse edges, say E and $\bar{E}$. Moreover, $f(E) \equiv E \cdot a$, where a is a path in G_{r-1} .*

We will use the following simple criterium to check, whether a given map between two 1-dimensional CW-complexes is a homotopy equivalence.

Proposition 4.4. *Let X, Y be finite connected 1-dimensional CW-complexes and let $f : X \rightarrow Y$ be a continuous map. Then f is a homotopy equivalence if and only if f induces an isomorphism between the fundamental groups $\pi_1(X, v)$ and $\pi_1(Y, f(v))$ for some vertex v of X .*

The following theorem says that in Theorem 4.3 we can provide $f(v) = v$ and choose p equal to the trivial path at v .

Theorem 4.5. *Let F be a free group of finite rank and α be an automorphism of F . Let n be the natural number and $f : \Gamma \rightarrow \Gamma$ be the relative train track from Theorem 4.3. Then one can algorithmically extend Γ to a finite connected graph Γ_1 , extend f to a relative train track $f_1 : \Gamma_1 \rightarrow \Gamma_1$, and indicate a vertex $v_1 \in \Gamma_1^0$ fixed by f_1 , and indicate an isomorphism $j : F \rightarrow \pi_1(\Gamma_1, v_1)$ such that the automorphism $j^{-1}\alpha^n j$ of the group $\pi_1(\Gamma_1, v_1)$ coincides with $(f_1)_*$. Moreover, one can provide the following property:*

(Pol) *Every polynomial stratum H_r consists of only two mutually inverse edges, say E and $\bar{E}$. Moreover, $f(E) \equiv E \cdot a$, where a is a path in G_{r-1} .*

Proof. Let $n, v, p, i : F \rightarrow \pi_1(\Gamma, v)$ be the number, the vertex, the path, and the isomorphism from Theorem 4.3, respectively. In particular, for every $w \in F$

we have

$$i(\alpha^n(w)) = [[p]]f_*(i(w))[[\bar{p}]]. \quad (4.1)$$

Let Γ_1 be the graph obtained from Γ by adding a new vertex v_1 and a new edge E connecting v_1 and $f(v)$. We extend the homotopy equivalence $f : \Gamma \rightarrow \Gamma$ to the map $f_1 : \Gamma_1 \rightarrow \Gamma_1$ by the rule $f_1(v_1) = v_1$ and $f_1(E) := Ef(p)$. Clearly, f_1 is a homotopy equivalence by Proposition 4.4. We define a maximal filtration for f_1 as extension of the maximal filtration for f with the help of the new top polynomial stratum consisting of the edges E and $\bar{E}$. Clearly, f_1 satisfies the property (Pol).

Finally, we define the isomorphism $j : F \rightarrow \pi_1(\Gamma_1, v_1)$ by the rule

$$j(w) := [[E]]f_*(i(w))[[\bar{E}]], \quad w \in F.$$

To complete the proof, we verify that the automorphism $j^{-1}\alpha^n j$ of the group $\pi_1(\Gamma_1, v_1)$ coincides with the induced automorphism $(f_1)_* : \pi_1(\Gamma_1, v_1) \rightarrow \pi_1(\Gamma_1, v_1)$. It suffices to check that $(f_1)_*(j(w)) = j(\alpha^n(w))$ for any $w \in F$. Using (4.1), we have

$$\begin{aligned} (f_1)_*(j(w)) &= (f_1)_*([[E]]f_*(i(w))[[\bar{E}]]]) = [[Ef(p)]](f^2)_*(i(w))[[f(\bar{p})\bar{E}]] = \\ &= [[E]]f_*(i(\alpha^n(w)))[[\bar{E}]] = j(\alpha^n(w)). \end{aligned}$$

□

For convenience, we reformulate Theorem 4.5 as follows.

Theorem 4.6. *Let F be a free group of finite rank and α be an automorphism of F . One can algorithmically compute a natural number n , construct a relative train track $f : (\Gamma, v) \rightarrow (\Gamma, v)$, and indicate an isomorphism $j : F \rightarrow \pi_1(\Gamma, v)$ such that the automorphism $j^{-1}\alpha^n j$ of the group $\pi_1(\Gamma, v)$ coincides with f_* . Moreover, one can provide the following property:*

(Pol) *Every polynomial stratum H_r consists of only two mutually inverse edges, say E and $\bar{E}$. Moreover, $f(E) \equiv E \cdot a$, where a is a path in G_{r-1} .*

Let F be a free group of finite rank with a fixed basis X . For any element $w \in F$ let $|w|$ be its word length with respect to X .

The following theorem was proven by P. Brinkmann in [13, Theorem 0.1] (an earlier inefficient version can be found in [14]).

Theorem 4.7. *There exists an algorithm which, given an automorphism α of a free group F of finite rank and given elements $u, v \in F$, verifies, whether there exists a natural N such that $\alpha^N(u) = v$. If such N exists, then the algorithm will compute N as well.*

Corollary 4.8. *Let Γ be a finite connected graph and $f : \Gamma \rightarrow \Gamma$ be a homotopy equivalence sending edges to edge-paths. Then for any two edge-paths ρ, τ in Γ , one can algorithmically verify, whether there exists a natural number k such that $f^k(\rho) = \tau$.*

Proof. First we reduce the problem to the case, where f fixes the endpoints of ρ . Let u_i and v_i be the initial and the terminal vertices of $\rho_i := f^i(\rho)$. Since f acts on the finite set $\Gamma^0 \times \Gamma^0$, there exist natural numbers r, n such that $(u_i, v_i) = (u_{i+n}, v_{i+n})$ for $i \geq r$.

First we check, whether $f^k(\rho) = \tau$ for $k < r$. If yes, we are done, if no we investigate the case $k \geq r$. Given such k , we can write $k = i + ln$ for some $l \geq 0$ and $r \leq i < r + n$. So, we have $f^k(\rho) = g^l(\rho_i)$, where $g := f^n$. Thus we have to investigate n problems: does there exist $l \geq 0$ such that $g^l(\rho_i) = \tau$, $r \leq i < r + n$. Note that g fixes the endpoints of ρ_i .

So, from the beginning, we may assume that f fixes the endpoints of ρ and $\alpha(\rho) = \alpha(\tau)$, and $\omega(\rho) = \omega(\tau)$.

Let Γ_1 be the graph obtained from Γ by adding a new vertex v and two new oriented edges: E_1 from v to $\alpha(\rho)$ and E_2 from v to $\omega(\rho)$. Let $f_1 : \Gamma_1 \rightarrow \Gamma_1$ be the extension of f which acts identically on E_1, E_2 . Clearly, f_1 is a homotopy equivalence which fixes v . Let $(f_1)_* : \pi_1(\Gamma_1, v) \rightarrow \pi_1(\Gamma_1, v)$ be the induced automorphism. We have

$$f^k(\rho) = \tau \iff f_1^k(E_1 \rho \bar{E}_2) = E_1 \tau \bar{E}_2 \iff (f_1)_*^k([E_1 \rho \bar{E}_2]) = [E_1 \tau \bar{E}_2].$$

Thus, the problem is solvable by Theorem 4.7. $\square$

Corollary 4.9. *Let Γ be a finite connected graph and $f : \Gamma \rightarrow \Gamma$ be a homotopy equivalence sending edges to edge-paths. Then for any two edge-paths ρ, τ in Γ , one can algorithmically verify, whether there exist natural numbers $k > s$ such that $f^k(\rho) = f^s(\tau)$.*

Proof. Let u_i and v_i be the initial and the terminal vertices of $f^i(\rho)$, $i \geq 0$, and let u'_j and v'_j be the initial and the terminal vertices of $f^j(\tau)$, $j \geq 0$. First we can decide, whether there exist i, j such that $(u_i, v_i) = (u'_j, v'_j)$. If such i, j don't exist, then the desired k, s don't exist. If such i, j exist, we can algorithmically find natural i, j, n with the following properties:

- 1) $(u_i, v_i) = (u'_j, v'_j)$;
- 2) $(u_i, v_i) = (u_{i+n}, v_{i+n})$ and n is minimal;
- 3) $i > j$;
- 4) $i - j$ is the minimal possible for 1)-3).

So, we reduce the problem to the following: does there exist $p \geq q \geq 0$ such that $f^{i+pn}(\rho) = f^{j+qn}(\tau)$? We set $\rho_1 := f^i(\rho)$, $\tau_1 := f^j(\tau)$, $g := f^n$. Then the endpoints

of ρ_1 and τ_1 coincide and are fixed by g . In this setting we have to decide, whether there exist $p \geq q \geq 0$ such that $g^p(\rho_1) = g^q(\tau_1)$.

We extend Γ to Γ' by adding an edge E from v_i to u_i and we extend g to $g' : \Gamma' \rightarrow \Gamma'$ by setting $g'|_\Gamma = g$ and $g(E) = E$. Then the problem is equivalent to the following: does there exist $p \geq q \geq 0$ such that $g'^p(\rho_1 E) = g'^q(\tau_1 E)$?

Since g' is a homotopy equivalence and $\rho_1 E$ and $\tau_1 E$ are loops based at the same point, and this point is fixed by g' , we have

$$g'^p(\rho_1 E) = g'^q(\tau_1 E) \iff g'^{p-q}(\rho_1 E) = \tau_1 E.$$

Thus, the problem can be reformulated as follows: does there exist $m \geq 0$ such that $g'^m(\rho_1 E) = \tau_1 E$? This can be decided by Theorem 4.7. $\square$

Theorem 4.10. *Let F be a free group of finite rank and let α be an automorphism of F of finite order. Then one can algorithmically compute a basis of $\text{Fix}(\alpha)$.*

Proof. Let H be the natural semidirect product of F and the finite cyclic group generated by α . In [29], Kalajdzewski gives an algorithm computing a finite generating set of $C_H(\alpha)$, the centralizer of α in H . Using the Reidemeister-Schreier method, one can easily compute a finite generating set of $\text{Fix}(\alpha) = C_H(\alpha) \cap F$. Finally, using Nielsen transformations or Stallings foldings, one can compute a basis of $\text{Fix}(\alpha)$. $\square$

Corollary 4.11. *Let α be an automorphism of a free group of finite rank and let n be a natural number. Suppose we know a basis of $\text{Fix}(\alpha^n)$. Then we can algorithmically compute a basis of $\text{Fix}(\alpha)$.*

Proof. The restriction of α to $\text{Fix}(\alpha^n)$ has finite order dividing n , and $\text{Fix}(\alpha)$ is a subgroup of $\text{Fix}(\alpha^n)$. Applying Theorem 4.10 to $F := \text{Fix}(\alpha^n)$ and $\alpha|_F$, we can compute a basis of $\text{Fix}(\alpha)$. $\square$

Thus, for computing a basis of $\text{Fix}(\alpha)$, it suffices to compute a basis of the group

$$\overline{\text{Fix}}(f) = \{[[\tau]] \in \pi_1(\Gamma, v) \mid f(\tau) = \tau\},$$

where Γ is the graph, v is the vertex and f is the relative train track from Theorem 4.6. In Section 9, we will show that we may assume that f satisfies some additional property (RTT-iv).

We need the following bounded cancelation lemma of Cooper, see [17], where it is credited to Grayson and Thurston.

Lemma 4.12. *Let Γ be a finite connected graph and $f : \Gamma \rightarrow \Gamma$ be a homotopy equivalence sending edges to edge-paths. Let τ_1, τ_2 be reduced paths in Γ such that $\omega(\tau_1) = \alpha(\tau_2)$ and the path $\tau_1 \tau_2$ is reduced. Then*

$$l([f(\tau_1 \tau_2)]) \geq l([f(\tau_1)]) + l([f(\tau_2)]) - 2C_*,$$

where $C_\star > 0$ is an algorithmically computable constant which depends only on f .

5. GRAPHS D_f AND $CoRe(C_f)$
FOR A HOMOTOPY EQUIVALENCE $f : \Gamma \rightarrow \Gamma$

In this section we recall some constructions and facts from [26, 37] and [16]. Let Γ be a finite connected graph with a distinguished vertex $v_\star$. Let $f : \Gamma \rightarrow \Gamma$ be a homotopy equivalence which maps vertices of Γ to vertices and which is locally injective on $\Gamma \setminus \Gamma^0$, and suppose that f fixes $v_\star$. We consider the group

$$\overline{\text{Fix}}(f) := \{[[p]] \in \pi_1(\Gamma, v_\star) \mid f(p) = p\}.$$

In papers [26, 37], the authors suggest a procedure for computation of a basis of $\overline{\text{Fix}}(f)$ with the help of a graph D_f . This procedure is not an algorithm in general case, since one cannot determine from the beginning, whether it terminates or not. We give a description of this procedure.

Definition of f -paths. An edge-path μ in Γ is called an f -path if the last point of μ coincides with the first point of $f(\mu)$.




Figure 2.
Nontrivial and trivial f -paths μ .

We note the following properties of f -paths:

- the trivial path at a vertex u of Γ , denoted $\mathbf{1}_u$, is an f -path if and only if u is fixed by f ;
- if μ is an f -path, then $[\mu]$ is also an f -path;
- if μ is an f -path and E is an edge in Γ such that $\alpha(E) = \alpha(\mu)$, then $\overline{E}\mu f(E)$ is also an f -path.

Definition of the graph D_f . The vertices of D_f are *reduced f -paths* in Γ . Let μ be a reduced f -path in Γ and let $E_1, \dots, E_n$ be all edges in Γ outgoing from $\alpha(\mu)$. Then we connect the vertex μ of D_f to the vertices $[\overline{E_1}\mu f(E_1)], \dots, [\overline{E_n}\mu f(E_n)]$ by edges with labels $E_1, \dots, E_n$, respectively, see Figure 3. The label of an edge-path in the graph D_f is the product of labels of consecutive edges of this path.

In general, the graph D_f can have infinitely many connected components and some of them can be infinite. For a vertex μ of D_f , let $D_f(\mu)$ be the component

of D_f containing μ . Lemma 5.3 says that $\pi_1(D(\mathbf{1}_{v_*}), \mathbf{1}_{v_*}) \cong \overline{\text{Fix}}(f)$. This was first proved in [26] with the help of preferable directions at vertices of D_f .

Preferable directions at vertices of D_f , dead and live vertices of D_f .

For a reduced nontrivial f -path μ in Γ , we set $\hat{f}(\mu) := [\overline{E}\mu f(E)]$, where E is the first edge of μ . Then μ and $\hat{f}(\mu)$ are vertices of the graph D_f connected by the edge with label E . The direction of this edge is called *preferable* at the vertex μ . We will put the symbol $\triangleright$ on this edge near the vertex μ .




Figure 3.

Note that only the vertices $\mathbf{1}_w$, where $w \in \Gamma^0$ and $f(w) = w$, do not admit a preferable direction. We call such vertices *dead* and all other *live*.

Observe that at all vertices of D_f , there is at most one outwardly directed edge.

Ordinary, repelling and attracting edges of D_f .

The following definition is illustrated by Figure 4.

Definition 5.1. Let e be an edge of D_f , let p, q be the initial and the terminal vertices of e , and let $E \in \Gamma^1$ be the label of e .

- (1) The edge e is called *ordinary* in D_f if one of the following holds:
 - (a) E is the first edge of the path p in Γ and $\overline{E}$ is not the first edge of the path q in Γ
 - (b) E is not the first edge of the path p in Γ and $\overline{E}$ is the first edge of the path q in Γ .

- (2) The edge e is called *repelling* in D_f if E is not the first edge of the path p in Γ and $\overline{E}$ is not the first edge of the path q in Γ .

A vertex of D_f is called *repelling* if it is a vertex of a repelling edge.

- (3) The edge e is called *attracting* in D_f if E is the first edge of the path p in Γ and $\overline{E}$ is the first edge of the path q in Γ .

An edge of D_f is called *exceptional* if it is attracting or repelling.




Figure 4. Different types of edges in D_f .

Proposition 5.2. (see [26, 37] and [16]) (a) *The repelling edges of D_f are in 1-1 correspondence with the occurrences of edges E in $f(E)$, where $E \in \Gamma^1$. More precisely, there exists a bijection of the type:*

$$f(E) \equiv \overline{u} \cdot E \cdot v \iff \left\{ \begin{array}{ll} \begin{array}{c} \text{Diagram 1: } \overline{u} \text{ (left arrow) } \xrightarrow{E} v \text{ (right arrow)} \\ \text{if } u \text{ and } v \text{ are nonempty,} \end{array} \\ \begin{array}{c} \text{Diagram 2: } \mathbf{1}_{\alpha(E)} \text{ (left arrow) } \xrightarrow{E} v \text{ (right arrow)} \\ \text{if } u \text{ is empty and } v \text{ not,} \end{array} \\ \begin{array}{c} \text{Diagram 3: } \overline{u} \text{ (left arrow) } \xrightarrow{E} \mathbf{1}_{\omega(E)} \text{ (left arrow)} \\ \text{if } v \text{ is empty and } u \text{ not,} \end{array} \\ \begin{array}{c} \text{Diagram 4: } \mathbf{1}_{\alpha(E)} \text{ (left arrow) } \xrightarrow{E} \mathbf{1}_{\omega(E)} \text{ (left arrow)} \\ \text{if } u \text{ and } v \text{ are empty.} \end{array} \end{array} \right.$$

(b) *The attracting edges of D_f are in 1-1 correspondence with the occurrences of edges $\overline{E}$ in $f(E)$, where $E \in \Gamma^1$. More precisely, there exists a bijection of the type:*

$$f(E) \equiv \overline{u} \cdot \overline{E} \cdot v \iff \begin{array}{c} \swarrow \quad \searrow \\ \bullet \quad \bullet \\ \nearrow \quad \nwarrow \end{array} \xrightarrow{E} \begin{array}{c} \nwarrow \quad \swarrow \\ \bullet \quad \bullet \\ \nearrow \quad \nwarrow \end{array}$$

$E \cdot u \qquad \qquad \qquad \overline{E} \cdot v$

(c) *There exist only finitely many exceptional edges in D_f and they can be algorithmically found.*

The following fundamental lemma was first proved by Goldstein and Turner in [26]. We reproduce here their nice proof for completeness.

Lemma 5.3. (see [26]) *The fundamental group of each component of D_f is finitely generated. Moreover, $\pi_1(D(\mathbf{1}_{v_*}), \mathbf{1}_{v_*}) \cong \overline{\text{Fix}}(f)$.*

Proof. Observe that a connected locally finite graph has finite rank if and only if the edges of this graph can be directed so that at all but a finite number of vertices, there is at most one outwardly directed edge.

Let D'_f be the graph obtained from D_f by removing all exceptional edges. Each component of D'_f has only ordinary edges, and these edges carry preferable directions which satisfy the above observation. Hence each component of D'_f has finite rank. Since there is only finitely many exceptional edges in D_f , each component of D_f has finite rank.

Let us prove the second claim in a little bit general context. Let μ be a vertex in D_f and suppose that c is a closed path in D_f based at μ . Let $\ell = E_1 E_2 \dots E_k$ be the label of c ; so all E_i are edges in Γ . Then

$$\mu, [\overline{E}_1 \mu f(E_1)], [\overline{E}_2 \overline{E}_1 \mu f(E_1) f(E_2)], \dots, [\overline{\ell} \mu f(\ell)]$$

are consecutive vertices of c and we have $[\overline{\ell} \mu f(\ell)] = \mu$.

So, ℓ is an edge-path in Γ based at $\alpha(\mu)$ and satisfying $[\mu f(\ell) \overline{\mu}] = [\ell]$. The correspondence $c \mapsto \ell$ induces the isomorphism

$$\pi_1(D_f(\mu), \mu) \cong \{[[p]] \in \pi_1(\Gamma, \alpha(\mu)) \mid [[\mu f(p) \overline{\mu}]] = [[p]]\}.$$

Setting $\mu := \mathbf{1}_{v_*}$, we obtain the second claim of the lemma. $\square$

Definition of a μ -subgraph of D_f . Let μ be a vertex in D_f . If μ is not a dead vertex, i.e. if $\mu \equiv E_1 E_2 \dots E_m$ for some edges $E_i \in \Gamma^1$, $m \geq 1$, we can pass from μ to the vertex $\widehat{f}(\mu) \equiv [E_2 \dots E_m f(E_1)]$ by using the direction which is preferable at μ .

The vertices of the μ -subgraph are the vertices $\mu_1, \mu_2, \dots$ of D_f such that $\mu_1 = \mu$ and $\mu_{i+1} = \widehat{f}(\mu_i)$ if the vertex μ_i is not dead, $i \geq 1$. The edges of the μ -subgraph are those which connect μ_i with μ_{i+1} and carry the preferable direction at μ_i .

Note that the μ -subgraph is finite if and only if starting from μ and moving along the preferable directions we will come to a dead vertex or to a vertex which we saw earlier. If the μ -subgraph is infinite, we call it a μ -ray. Thus, any μ -subgraph is one of the following four types:




Figure 5.

Let μ and τ be two vertices of D_f . Clearly, if the μ -subgraph and the τ -subgraph intersect, then they differ only by their finite “initial subsegments”. The vertex μ_i from this intersection with minimal i is called the *intersection point* of these subgraphs.

Note the following properties of μ -subgraphs:

- if μ_0 is a vertex of a μ -subgraph, then the μ_0 -subgraph is contained in the μ -subgraph;
- if μ_0 is a vertex of a μ -subgraph and τ_0 is a vertex of a τ -subgraph, then the μ -subgraph and the τ -subgraph intersect if and only if the μ_0 -subgraph and the τ_0 -subgraph intersect;
- a μ -ray does not intersect a finite τ -subgraph.

Definitions of the graphs C_f and $CoRe(C_f)$. A component of D_f is called *repelling* if it contains at least one repelling edge. Let $C_1, \dots, C_n$ be all repelling components of D_f . For each C_i , let $CoRe(C_i)$ be a core of C_i which contains all repelling edges of C_i . We set $C_f := \bigcup_{i=1}^n C_i$ and $CoRe(C_f) := \bigcup_{i=1}^n CoRe(C_i)$.

Below we show how to compute a basis of the group $\pi_1(D_f(\mathbf{1}_{v_*}), \mathbf{1}_{v_*})$ if we know how to construct the graph $CoRe(C_f)$.

Lemma 5.4. *Let $\mathbf{1}_u$ be a dead vertex of D_f . If the component $D_f(\mathbf{1}_u)$ is non-contractible, then it lies in C_f .*

Proof. Suppose that $D_f(\mathbf{1}_u)$ is non-contractible. Then there exists an edge-path $p = E_1 E_2 \dots E_k$ in $D_f(\mathbf{1}_u)$ such that $\omega(E_k) = \mathbf{1}_u$, the edges of p are distinct, and $\alpha(E_1) = \omega(E_s)$ for some $1 \leq s \leq k$. We show that p contains a repelling edge. Suppose not, then the direction of E_k is preferable at $\alpha(E_k)$. By induction, the direction of E_i is preferable at the point $\alpha(E_i)$ for every $i = 1, \dots, k$ (see Figure 6). In particular, $\alpha(E_1) \neq \mathbf{1}_u$, and hence $k > s$. Then there are two preferable directions at $\alpha(E_1)$, namely the direction of E_1 and the direction of E_{s+1} , a contradiction. Thus, p must contain a repelling edge. $\square$




Figure 6.

Lemma 5.5. *Let $\mathbf{1}_u$ be a dead vertex of D_f . The vertex $\mathbf{1}_u$ lies in C_f if and only if it lies in a μ -subgraph for some repelling vertex μ .*

Proof. Suppose that $\mathbf{1}_u$ lies in C_f . Then there exists a shortest path $E_1 E_2 \dots E_k$, where the edge E_1 is repelling and $\omega(E_k) = \mathbf{1}_u$. If $k = 1$, then $\mathbf{1}_u$ is repelling as a vertex of a repelling edge, and we are done. If $k \geq 2$, the edges $E_2, \dots, E_k$ are not repelling. In particular, the preferable direction at $\alpha(E_k)$ coincides with the direction of E_k . By induction one can prove that the preferable direction at $\alpha(E_i)$ coincides with the direction of E_i for $i \geq 2$. Then $\mathbf{1}_u$ lies in the μ -subgraph for $\mu := \omega(E_1)$ and the vertex μ is repelling. The converse claim is clear. $\square$

Proposition 5.6. *Suppose we can algorithmically construct $\text{CoRe}(C_f)$ and decide whether the vertex $\mathbf{1}_{v_*}$ lies in the μ -subgraph for some repelling vertex μ . Then we can compute a basis of $\pi_1(D_f(\mathbf{1}_{v_*}), \mathbf{1}_{v_*})$.*

Proof. By assumption and Lemma 5.5, we can algorithmically verify, whether $\mathbf{1}_{v_*}$ lies in C_f . If $\mathbf{1}_{v_*} \notin C_f$, then $D_f(\mathbf{1}_{v_*})$ is contractible by Lemma 5.4.

If $\mathbf{1}_{v_*} \in C_f$, then $D_f(\mathbf{1}_{v_*})$ is a component of C_f . By Lemma 5.5, we can find a repelling vertex μ such that the μ -subgraph terminates at $\mathbf{1}_{v_*}$. Obviously, the μ -subgraph lies in $D_f(\mathbf{1}_{v_*})$.

Since $CoRe(C_f)$ is supposed to be constructible, we can find the component of $CoRe(C_f)$ containing μ . Let Δ be the union of this component and the μ -subgraph. Then Δ is a core of $D_f(\mathbf{1}_{v_*})$ containing $\mathbf{1}_{v_*}$. In particular, we can compute a basis of $\pi_1(D_f(\mathbf{1}_{v_*}), \mathbf{1}_{v_*})$. $\square$

To construct the graph $CoRe(C_f)$, it suffices to do the following:

- (1) Find all repelling edges of D_f .
- (2) For each live repelling vertex μ determine, whether the μ -subgraph is finite or not.
- (3) Compute all elements of all finite μ -subgraphs from (2).
- (4) For each two repelling vertices μ and τ with infinite μ - and τ -subgraphs determine, whether these subgraphs intersect.
- (5) If the μ -subgraph and the τ -subgraph from (4) intersect, find their first intersection point and compute their initial segments up to this point.

To convert this procedure to an algorithm, we shall construct algorithms for steps (2) and (4). In papers [16] and [37] these algorithms are given only in some special cases (for positive and for irreducible automorphisms represented by train tracks for which each fixed point is a vertex). The main idea in these papers is to use an *inverse preferred direction* at vertices in the graph D_f . This direction can be constructed algorithmically (in general case) with the help of a homotopic inverse to f . It determines its own repelling edges, repelling and dead vertices; they can be algorithmically found.

Inverse preferred directions in D_f . We will realize the following plan. First we will define a map $g : \Gamma \rightarrow \Gamma$ which is a homotopy inverse to $f : \Gamma \rightarrow \Gamma$. Then we show that there is a label preserving graph map $\Phi : D_f \rightarrow D_g$. Finally we define the inverse preferred directions for vertices in D_f by pulling back the preferred directions in D_g by Φ . This idea is due to Turner [37], and has sources in the paper of Cohen and Lustig [16]. Note that in [37], the map Φ is claimed to be locally injective (see Proposition in Section 3 there), and we claim that Φ is an isomorphism.

For the given homotopy equivalence $f : \Gamma \rightarrow \Gamma$, we can algorithmically find a homotopy equivalence $g : \Gamma \rightarrow \Gamma$ such that g maps vertices of Γ to vertices, edges to edge-paths, and the maps $h := g \circ f$ and $f \circ g$ are homotopic to the identity on Γ . Let $H : \Gamma \times [0, 1] \rightarrow \Gamma$ be a homotopy from the identity id to h . For each vertex u in Γ , let p_u be the path from u to $h(u)$ determined by the homotopy H : namely $p_u(t) = H(u, t)$, $t \in [0, 1]$.

First we define a map Φ from the set of vertices of D_f to the set of vertices of D_g . Let μ be a vertex in D_f . We consider μ as a reduced f -path in Γ and let u be the initial vertex of μ . Then we set $\Phi(\mu) = [p_u g(\bar{\mu})]$. Clearly, $\Phi(\mu)$ is a reduced g -path in Γ . Hence $\Phi(\mu)$ can be considered as a vertex in D_g .

Lemma 5.7. *The map Φ can be continued to a graph map $\Phi : D_f \rightarrow D_g$ preserving the labels of edges.*

Proof. Let μ and μ_1 be two vertices in D_f connected by an edge with label E , i.e. $\mu_1 = [\overline{E}\mu f(E)]$. We must show that $\Phi(\mu)$ and $\Phi(\mu_1)$ are connected by an edge with the label E , i.e. $\Phi(\mu_1) = [\overline{E}\Phi(\mu)g(E)]$. Let u and w be the initial and the terminal vertices of E . Then u and w are the initial vertices of μ and μ_1 , respectively. We have

$$\Phi(\mu_1) = [p_w g(f(\overline{E})\overline{\mu}E)] = [p_w h(\overline{E})g(\overline{\mu})g(E)] = [\overline{E}p_u g(\overline{\mu})g(E)] = [\overline{E}\Phi(\mu)g(E)].$$

Here we use the fact that $h(E)$ is homotopic to $\overline{p}_u E p_w$ with respect to the endpoints. The latter follows from the continuity of the map $H|(E \times [0, 1])$. $\square$

Remark 5.8. Let μ be a vertex in D_f . Then the following holds:

- 1) The f -path μ and the g -path $\Phi(\mu)$ have the same initial vertices in Γ .
- 2) Let $E_1, \dots, E_n$ be edges outgoing from $\alpha(\mu)$ in Γ . Then the vertices μ and $\Phi(\mu)$ of the graphs D_f and D_g have degree n and the labels of edges outgoing from each of these vertices are $E_1, \dots, E_n$.
- 3) Φ maps the star of the vertex μ to the star of the vertex $\Phi(\mu)$ bijectively and label preserving.

Proposition 5.9. *The map $\Phi : D_f \rightarrow D_g$ is an isomorphism of graphs.*

Proof. By Remark 5.8. 3), it suffices to show that Φ is bijective on vertices.

First we show that Φ is injective on vertices. Let μ_1, μ_2 be two different vertices of D_f . If the f -paths μ_1 and μ_2 have different initial vertices in Γ , then, by Remark 5.8. 1), the g -paths $\Phi(\mu_1)$ and $\Phi(\mu_2)$ have different initial vertices in Γ too, hence $\Phi(\mu_1) \neq \Phi(\mu_2)$.

Suppose that the initial vertices of the f -paths μ_1 and μ_2 coincide and equal to u . Then their terminal vertices also coincide and equal to $f(u)$. Since the f -paths μ_1, μ_2 are reduced, $\mu_1 \neq \mu_2$, and g is a homotopy equivalence, we have $[g(\mu_1)] \neq [g(\mu_2)]$, hence $\Phi(\mu_1) = [p_u g(\overline{\mu}_1)] \neq [p_u g(\overline{\mu}_2)] = \Phi(\mu_2)$.

Now we show that Φ is surjective on vertices. Let τ be a vertex in D_g , i.e. τ is a reduced g -path in Γ . Let u be the initial vertex of the path τ . We will find a reduced f -path μ in Γ such that $\Phi(\mu) = \tau$. Let μ_1 be an arbitrary path in Γ from u to $f(u)$. Then the paths τ and $p_u g(\overline{\mu}_1)$ have the same endpoints, so $\overline{\tau} p_u g(\overline{\mu}_1)$ is a loop based at $g(u)$. Hence, there exists a loop σ in Γ based at u such that $g(\sigma) = \overline{\tau} p_u g(\overline{\mu}_1)$. We set $\mu := [\sigma \mu_1]$. Then $\Phi(\mu) = [p_u g(\overline{\mu})] = [p_u g(\overline{\mu}_1)g(\overline{\sigma})] = \tau$. $\square$

Definition 5.10. The *inverse preferred direction* at a vertex μ in D_f is the preimage of the preferred direction at the vertex $\Phi(\mu)$ in D_g under Φ .

We formulate this more detailed. Recall that $\Phi(\mu) = [p_u g(\bar{\mu})]$, where u is the initial vertex of the f -path μ . First suppose that the f -path $\Phi(\mu)$ is nontrivial and let E be the first edge of this path. Then the inverse preferred direction at the vertex μ of D_f is the direction of the edge of D_f which starts at μ and has the label E .

If the f -path $\Phi(\mu)$ is trivial in Γ , the inverse preferred direction at μ in D_f is not defined.

Proposition 5.11. *The inverse preferred direction is defined at almost all vertices of D_f .*

Proof. It suffices to show that for each vertex u in Γ , there exists at most one reduced f -path μ with $\alpha(\mu) = u$ and trivial $\Phi(\mu)$. Suppose that there are two such paths μ_1, μ_2 . Then we have $[p_u g(\bar{\mu}_1)] = [p_u g(\bar{\mu}_2)]$, and hence $[g(\mu_1)] = [g(\mu_2)]$. Since μ_1 and μ_2 are f -paths with the same initial vertex u and the same terminal vertex $f(u)$, and g is a homotopy equivalence, we have $\mu_1 = \mu_2$. $\square$

Definition 5.12. Preimages, with respect to Φ , of repelling edges, repelling vertices and dead vertices of D_g are called *inv-repelling edges*, *inv-repelling vertices* and *inv-dead vertices* of D_f .

Note that there are only finitely many inv-repelling edges and inv-repelling and inv-dead vertices in D_f , and they can be algorithmically found.

Definition 5.13. A vertex of D_f is called *normal* if the preferred and the inverse preferred directions at this vertex exist and do not coincide.

The following propositions are contained in Claim b) in the proof of Theorem A in [37]. This claim was inspired by Lemma (4.8) and Proposition (4.10) from [16].

Proposition 5.14. *Suppose that R is a μ -ray in D_f . Then the preferred direction on all but finitely many edges in R is opposite to the inverse preferred direction. In particular, almost all vertices in R are normal.*

Proposition 5.15. *Let R_1 and R_2 be a μ_1 -ray and a μ_2 -ray in D_f , respectively. Suppose that they do not contain inv-repelling vertices and that their initial vertices μ_1 and μ_2 are normal. Then R_1 and R_2 are either disjoint or one is contained in the other.*

How to convert the procedure (1)-(5) for construction of the graph $\text{CoRe}(C_f)$ into an algorithm

As it was observed, it suffices to find algorithms for steps (2) and (4).

Using Propositions 5.14 and 5.15, Step (4) can be replaced by the following three steps.

- (4.1) For each repelling vertex μ whose μ -subgraph is a ray, find in this μ -ray a vertex μ' such that the μ' -ray does not contain inv-repelling vertices.
- (4.2) Find a normal vertex μ'' in the μ' -ray.
- (4.3) For every two repelling vertices μ and τ , whose μ - and τ -subgraphs are rays, verify whether τ'' is contained in the μ'' -ray or μ'' is contained in the τ'' -ray.

Step (4.2) can be done algorithmically because of Proposition 5.14: we move along the μ' -ray and await the appearance of the first normal vertex. Steps (4.1) and (4.3) can be done if we find an algorithm for the following problem.

Problem 1. Given two vertices μ and τ of the graph D_f , verify whether τ is contained in the μ -subgraph.

Indeed, for Step (4.1) we first find all inv-repelling vertices in D_f . Then we detect those of them which lie in the μ -ray. Let I be the minimal initial segment of the μ -ray which contains all these vertices. We can take μ' equal to the first vertex in the μ -ray which lies outside I . Step (4.3) is a partial case of Problem 1.

Step (2) can be done if we find an algorithm for the following problem:

Problem 2. Given a live vertex μ of the graph D_f , determine whether the μ -subgraph is finite or not.

Thus, to construct $CoRe(C_f)$ algorithmically, it suffices to find algorithms for Problems 1 and 2. Moreover, using an algorithm for Problem 1, we can decide whether the vertex $\mathbf{1}_{v_*}$ lies in a μ -subgraph for some repelling vertex μ . Then, by Proposition 5.6 we can compute a basis of $\pi_1(D_f(\mathbf{1}_{v_*}), \mathbf{1}_{v_*})$. Lemma 5.3 identifies this group with $\overline{\text{Fix}}(f)$.

In Section 15 we will present algorithms for Problems 1 and 2 in case, where $f : \Gamma \rightarrow \Gamma$ is a subdivided relative train track satisfying the condition (Pol). As explained in Section 3, this will provide an algorithm for computing $\text{Fix}(\alpha)$.

6. CANCELLATION POINTS IN THE PRODUCT OF TWO PATHS

Let H_r be an exponential stratum of a relative train track $f : \Gamma \rightarrow \Gamma$.

Definition 6.1. Let τ be a reduced edge-path in G_r and y be an occurrence of a vertex in τ different from the endpoints of τ . We write $\tau \equiv \bar{p}_0 \cdot q_0$, where q_0 is the final subpath of τ starting at y . Let A be the first edge of p_0 and B be the first edge of q_0 . The vertex y is called an *r-cancellation point* in τ if (A, B) is an illegal *r*-turn.

Let $\mu, \tau \subset G_r$ be reduced edge-paths, such that $\omega(\mu) = \alpha(\tau)$. Let $y_k, \dots, y_1$ and $z_1, \dots, z_s$ be the consequent r -cancellation points in μ and τ , respectively. We denote by $I(\bar{\mu}, \tau)$ the largest common subpath of $\bar{\mu}$ and τ , and let t be the terminal vertex of $I(\bar{\mu}, \tau)$. Then we have one of the following four cases for the path $\mu\tau$ (see Figure 7).




Figure 7.

(1) Suppose that $t \notin \{y_1, \dots, y_k, z_1, \dots, z_s\}$ and that $y_1 = z_1, \dots, y_l = z_l$, are all r -cancellation points in $I(\bar{\mu}, \tau)$. Then all r -cancellation points in $[\mu\tau]$ are $y_k, \dots, y_{l+1}, z_{l+1}, \dots, z_s$ and, possibly, t (see Figure 7 (1)).

(2) Suppose that $t = y_l = z_l$ for some l . Then t is an r -cancellation point in $[\mu\tau]$. This follows from the fact that the turns at t in μ and τ are illegal and have a common edge. So, all r -cancellation points in $[\mu\tau]$ are $y_k, \dots, y_{l+1}, t, z_{l+1}, \dots, z_s$ (see Figure 7 (2)).

(3) Suppose that $t = y_l$ for some l and that $t \notin \{z_1, \dots, z_s\}$. Then $y_1 = z_1, \dots, y_{l-1} = z_{l-1}$ are all r -cancellation points in the path $I(\bar{\mu}, \tau)$. Moreover, t is not an r -cancellation point in $[\mu\tau]$. This follows from the fact that the turn at t in μ is illegal, while the turn at t in τ is legal and these turns have a common edge. So, all r -cancellation points in $[\mu\tau]$ are $y_k, \dots, y_{l+1}, z_l, \dots, z_s$ (see Figure 7 (3)).

(4) Suppose that $t = z_l$ for some l and that $t \notin \{y_1, \dots, y_k\}$ (see Figure 7 (4)). This case is analogous to Case (3). In particular, t is not an r -cancellation point in $[\mu\tau]$.

Thus, the maximal possible number of r -cancellation points in $[\mu\tau]$ is $k + s + 1$ (it can be achieved only in Case (1) for $l = 0$). If at least one of the r -cancellation points in μ and τ lies in $I(\bar{\mu}, \tau)$, then the number of r -cancellation points in $[\mu\tau]$ is strictly less than $k + s$.

7. NON-DELETABLE CANCELLATION POINTS IN EXPONENTIALLY GROWING PATHS

Let Γ be a finite connected graph, $f : \Gamma \rightarrow \Gamma$ be a relative train track and let $\emptyset = G_0 \subset \dots \subset G_N = \Gamma$ be the maximal filtration in Γ . In this section, we fix a natural number $0 < r \leq N$ and assume that $H_r := \text{cl}(G_r \setminus G_{r-1})$ is an exponential stratum. Let λ be the eigenvalue corresponding to H_r .

We introduce some notations. Let $\tau \equiv E_1 \dots E_n$ be a reduced edge-path in the subgraph G_r . If for some $1 \leq i \leq n - 1$ the turn $(\overline{E_i}, E_{i+1})$ is illegal, then we say that τ contains the illegal turn $(\overline{E_i}, E_{i+1})$ at the vertex $\omega(E_1 \dots E_i)$. Using Properties (RTT-i)–(RTT-iii), one can show that the number of illegal r -turns in $[f(\tau)]$ does not exceed the number of illegal r -turns in τ .

We show how to decide whether the number of illegal r -turns in the sequence $\tau, [f(\tau)], [f^2(\tau)], \dots$ is constant. Figure 8 aims to show that illegal r -turns in τ can behave differently under applying f .




Figure 8.

The bold vertices are vertices of illegal turns. Picture (1) shows two illegal turns in τ which yield two illegal turns in $[f^2(\tau)]$. Picture (2) shows that the number of illegal turns can be decreased in two steps.

Definition 7.1. Let p_0, q_0 be reduced edge-paths in Γ with the same initial vertex. By $I(p_0, q_0)$ we denote the largest common initial segment of p_0 and q_0 . Then $p_0 \equiv I(p_0, q_0) \cdot p_1$ and $q_0 \equiv I(p_0, q_0) \cdot q_1$ for some reduced paths p_1, q_1 . We denote by $\Lambda(p_0, q_0)$ the ordered pair of paths (p_1, q_1) .

Let τ be a reduced edge-path in the subgraph G_r and let y be a vertex of τ . We write $\tau \equiv \bar{p}_0 \cdot q_0$, where q_0 is the final subpath of τ starting at y . Set

$$(p_k, q_k) \equiv \Lambda([f^k(p_0)], [f^k(q_0)]) \quad \text{and} \quad \alpha_k \equiv I([f(p_{k-1})], [f(q_{k-1})])$$

for $k \geq 1$ (see Figure 9). Then $[f^k(\tau)] \equiv \bar{p}_k \cdot q_k$. The occurrence $y^k := \alpha(p_k) = \alpha(q_k)$ in $[f^k(\tau)]$ is called the k -successor of y .




Figure 9.

Definition 7.2. Let τ be a reduced edge-path in G_r with a unique r -cancellation point y (see Definition 6.1). We say that y is *non-deletable* in τ if for every $k \geq 1$ the k -successor y^k is an r -cancellation point in $[f^k(\tau)]$. We say that y is *deletable* in τ if this does not hold.

Observe that for every $k \geq 1$ the following is valid: the point y^k is a non-deletable r -cancellation point in $[f^k(\tau)]$ if and only if y is a non-deletable r -cancellation point in τ .

Definition 7.3. Let τ be a reduced edge-path in G_r with a unique r -cancellation point y and suppose that y is non-deletable. The point y divides τ into two parts, say $\bar{p}_0$ and q_0 , so $\tau \equiv \bar{p}_0 \cdot q_0$. A pair (p'_0, q'_0) , where p'_0 and q'_0 are initial edge-subpaths of p_0 and q_0 , respectively, is called *responsible for r -cancellations in τ* if $I([f^i(p_0)], [f^i(q_0)]) \equiv I([f^i(p'_0)], [f^i(q'_0)])$ for every $i \geq 0$. Let $((p_0)_{\min}, (q_0)_{\min})$ be the pair which is minimal, with respect to the sum of l -lengths of components, among all pairs responsible for r -cancellations in τ .

By this definition, every edge of each of the segments $(p_0)_{\min}$ and $(q_0)_{\min}$ except the last one can be moved by appropriate f^i into the segment $I_i := I([f^i(p_0)], [f^i(q_0)])$.

Lemma 7.4. *Let τ be a reduced edge-path in G_r with a unique r -cancellation point y and suppose that y is non-deletable. With the above notations the following statements hold:*

- 1) *The first and the last edges of the paths $(p_0)_{\min}$ and $(q_0)_{\min}$ belong to H_r .*
- 2) *There exists a computable real number L_{critical} (depending only on f), such that $L_r((p_0)_{\min}) < L_{\text{critical}}$ and $L_r((q_0)_{\min}) < L_{\text{critical}}$. In particular, the number of r -edges in $(p_0)_{\min}$ and in $(q_0)_{\min}$ is bounded from above by a computable natural number n_{critical} (depending only on f).*

Proof. 1) The initial edges of $(p_0)_{\min}$ and $(q_0)_{\min}$ form an r -turn (this follows from the definition of an r -cancellation point), so they belong to H_r . We prove that the terminal edges of $(p_0)_{\min}$ and $(q_0)_{\min}$ belong to H_r . Suppose the contrary: say the last edge of $(p_0)_{\min}$ lies in G_{r-1} . Then we can write $(p_0)_{\min} \equiv ab$, where a begins and ends on r -edges and b is a nontrivial edge-path in G_{r-1} . We have

$$[f^k((p_0)_{\min})] \equiv [f^k(a)] \cdot [f^k(b)].$$

Since $(p_0)_{\min}$ is minimal, there exists $k_0 \geq 1$ such that $[f^k(a)]$ is contained in

$$I_k \equiv I([f^k(p_0)], [f^k(q_0)]) \equiv I([f^k((p_0)_{\min})], [f^k((q_0)_{\min})])$$

for every $k \geq k_0$. In particular, the H_r -parts of $[f^k(a)]$ and I_k coincide for every $k \geq k_0$. This and the equation $I_{k+1} \equiv [f(I_k)]\alpha_{k+1}$ imply that α_{k+1} lies in G_{r-1} for every $k \geq k_0$. But since y is non-deletable, there exists $t = t(k) \geq 1$ such that α_{k+t} is nonempty, hence it begins on an r -edge, a contradiction.

2) We set $L_{\text{critical}} := M + L$, where

$$M := \max\{L_r(E) \mid E \in H_r^1\}, \quad L := \frac{C_\star}{\lambda - 1} M,$$

and $C_\star$ is the Cooper constant from Lemma 4.12.

With the above notations we have $\tau \equiv \bar{p}_0 q_0$. Suppose that $L_r((p_0)_{\min}) \geq L_{\text{critical}}$. Then there exists a decomposition $p_0 \equiv w_0 u$, where w_0 is the initial edge-subpath of p_0 of minimal length which ends on an r -edge and satisfies

$$L_r(w_0) \geq L.$$

By minimality, we have

$$L_r(w_0 \text{ without the last edge}) < L,$$

and hence $L_r(w_0) < L_{\text{critical}}$. To get a contradiction, it suffices to show that $(p_0)_{\min}$ is an initial segment of w_0 .

Since p_0 is r -legal, w_0 is r -legal as well, and hence

$$L_r([f(w_0)]) \geq \lambda L.$$

Since $\tau \equiv \bar{p}_0 q_0$ is reduced, in the product $[f(\bar{p}_0)][f(q_0)]$ at most $C_\star$ edges can be canceled at the end of $[f(\bar{p}_0)]$. Canceling, we have $[f(\tau)] \equiv \bar{p}_1 q_1$, where $p_1 \equiv w_1[f(u)]$ for some final segment w_1 of $[f(w_0)]$ with

$$L_r(w_1) \geq L_r([f(w_0)]) - C_\star \cdot M \geq L.$$

Moreover, w_1 as well as $[f(w_0)]$ ends on an r -edge.

Continuing, we obtain $[f^k(\tau)] \equiv \bar{p}_k q_k$, where $p_k \equiv w_k[f^k(u)]$ for some w_k which ends on an r -edge and satisfies

$$L_r(w_k) \geq L.$$

In particular, no point of u can be moved into $I([f^k(p_0)], [f^k(q_0)])$ with the help of f^k . Therefore, $(p_0)_{\min}$ is an initial segment of w_0 , a contradiction.

The number of r -edges in $(p_0)_{\min}$ and in $(q_0)_{\min}$ does not exceed L_{critical}/M' , where $M' := \min\{L_r(E) \mid E \in H_r^1\}$. So, we can set $n_{\text{critical}} := \lceil L_{\text{critical}}/M' \rceil$. $\square$

Proposition 7.5. *Let τ be a reduced edge-path in G_r . Suppose that each of the paths τ , $[f(\tau)]$, $[f^2(\tau)]$, $\dots$ contains a unique illegal r -turn. Then there exist integer numbers $T \geq 0$ and $p \geq 1$ such that*

$$((p_{T+j})_{\min}, (q_{T+j})_{\min}) \equiv ((p_{T+p+j})_{\min}, (q_{T+p+j})_{\min})$$

for all $j \geq 0$. In particular,

$$\alpha_{T+j} \equiv \alpha_{T+p+j}$$

for all $j \geq 1$. Moreover, T and p are computable and depend only on f , but not on τ .

Proof. Let $(p_0)_{\min} \equiv a_1 b_1 \dots a_s b_s a_{s+1}$ and $(q_0)_{\min} \equiv a'_1 b'_1 \dots a'_t b'_t a'_{t+1}$, where all a_i and a'_i are nontrivial and lie in H_r and all b_i and b'_i are nontrivial and lie in G_{r-1} . We use notation $I_k := I([f^k(p_0)], [f^k(q_0)])$. Recall that $[f^k(p_0)] \equiv I_k p_k$ and $[f^k(q_0)] \equiv I_k q_k$. We set $z_k := \alpha(p_k)$.

Claim 1. There exists a computable natural number k_0 (depending only on f) such that for all $k \geq k_0$ and every b_i , $i = 1, \dots, s$, one of the following holds:

- 1) $[f^k(b_i)]$ is a subsegment of I_k . In this case we say that the segment b_i can be moved into I_k in k steps.
- 2) $[f^k(b_i)]$ is a subsegment of p_k and the l -distance from $z_k = \alpha(p_k)$ to $\alpha([f^k(b_i)])$ along p_k is at most $\|f\|^{k_0}$. (We stress that the last number does not depend on k .)

The same alternative holds for every b'_j , $j = 1, \dots, t$.

Proof. Let k_0 be the minimal integer such that

$$\lambda^{k_0} \min\{L_r(E) \mid E \in H_r^1\} > L_{\text{critical}} \quad (7.1)$$

and let $k \geq k_0$. We fix $i \in \{1, \dots, s\}$. Let e be the last edge of a_i , so e is an r -edge and $\omega(e) = \alpha(b_i)$. Let E be the last edge of $[f^{k-k_0}(e)]$, so E is an r -edge and $\omega([f^{k_0}(E)]) = \alpha([f^k(b_i)])$. Note that $[f^{k_0}(E)]$ is a subsegment of $[f^k(e)]$, which is a subsegment of $[f^k(p_0)] \equiv I_k p_k$.

Case 1. Suppose that $[f^{k_0}(E)]$ lies in I_k . Then $\alpha([f^k(b_i)])$ lies in I_k . This implies that the whole segment $[f^k(b_i)]$ lies in I_k , otherwise it would cover the first edge of p_k that is impossible, since this edge lies in H_r but $[f^k(b_i)]$ lies in G_{r-1} . So, in this case the statement 1) holds.

Case 2. Suppose that $[f^{k_0}(E)]$ does not lie in I_k . Since $L_r([f^{k_0}(E)]) > L_{\text{critical}}$ by (7.1), the segment $[f^{k_0}(E)]$ does not lie in p_k as well; otherwise $[f^{k_0}(E)]$ could never be moved into appropriate I_l . Therefore $\alpha([f^{k_0}(E)])$ lies in I_k and $\omega([f^{k_0}(E)])$ lies in p_k . Then $[f^{k_0}(E)]$ covers z_k . So, the l -distance from z_k to $\omega([f^{k_0}(E)]) = \alpha([f^k(b_i)])$ along p_k is at most $\|f\|^{k_0}$. Then the statement 2) holds. $\square$

Claim 2. There exists a computable natural number k_1 (depending only on f) such that at least one of the segments b_s, b'_t can be moved into I_{k_1} in k_1 steps.

Proof. We prove that this claim holds for $k_1 := k_0 + (N^2 + 1)$, where N is the number of the edge-paths in G_r with l -length at most $\|f\|^{k_0}$. Suppose the contrary. Then, by Claim 1 the following is satisfied for every $k = k_0, \dots, k_1$: the segment of p_k from $\alpha(p_k)$ to $\alpha([f^k(b_s)])$ and the segment of q_k from $\alpha(q_k)$ to $\alpha([f^k(b'_t)])$ have l -lengths at most $\|f\|^{k_0}$. But then these pairs of segments will repeat, hence, for every natural j , the segments b_s and b'_t cannot be moved into I_j . This contradicts the minimality of $(p_0)_{\min}$ and $(q_0)_{\min}$. $\square$

Claim 3. There exists a computable natural number k_2 (depending only on f) such that both segments b_s, b'_t can be moved into I_{k_2} in k_2 steps.

Proof. By Claim 2, we may assume that b'_t can be moved into I_{k_1} in k_1 steps. Then the initial vertex of a'_{t+1} (but not the whole segment a'_{t+1}) can be moved into I_{k_1} in k_1 steps. This means that $(q_{k_1})_{\min}$ is a subpath of $[f^{k_1}(a'_{t+1})]$. Let N_1 be the maximum of the number of edge-subpaths in $[f^{k_1}(\ell)]$ over all paths $\ell \subset H_r$ with $L_r(\ell) < L_{\text{critical}}$. Then there is at most N_1 possibilities for $(q_{k_1})_{\min}$. We prove that we can take $k_2 := k_1(N \cdot N_1 + 1)$.

If for some $i = 1, \dots, (N \cdot N_1 + 1)$, the path b_s can be moved into $I_{k_1 i}$ in $k_1 i$ steps, we are done. If not, then by Claim 1, the initial segment of $(p_{k_1 i})_{\min}$ from $z_{k_1 i}$ to the beginning of $[f^{k_1 i}(b_s)]$ has l -length at most $\|f\|^{k_0}$. There are N possibilities for such segments and there are N_1 possibilities for $(q_{k_1 i})_{\min}$, $i = 1, \dots, (N \cdot N_1 + 1)$. Hence, we obtain a repeating pair consisting of the initial segment of $(p_{k_1 i})_{\min}$ from

$z_{k_1 i}$ to the beginning of $[f^{k_1 i}(b_s)]$ and of $(q_{k_1 i})_{\min}$. This implies that b_s cannot be moved into I_j for every j , a contradiction. $\square$

Claim 4. We can find algorithmically the desired T and p .

Proof. By Claim 3, the initial vertices of the segments $[f^{k_2}(a_{s+1})]$ and $[f^{k_2}(a'_{t+1})]$ lie in I_{k_2} . Since their terminal vertices lie in p_{k_2} and in q_{k_2} respectively, $(p_{k_2})_{\min}$ and $(q_{k_2})_{\min}$ are some subsegments of $[f^{k_2}(a_{s+1})]$ and $[f^{k_2}(a'_{t+1})]$. Since $a_{s+1}, a'_{t+1} \subset H_r$ are parts of $(p_0)_{\min}$ and $(q_0)_{\min}$, respectively, we have $l(a_{s+1}) \leq n_{\text{critical}}$ and $l(a'_{t+1}) \leq n_{\text{critical}}$ by Lemma 7.4. Therefore, there is only finitely many possibilities for $(p_{k_2})_{\min}$ and $(q_{k_2})_{\min}$ (this number is computable and depends only on f). Thus, we can compute a universal constant N_2 (depending only on f) for which there exist numbers $1 < i < i + d < N_2$ such that the pairs $((p_i)_{\min}, (q_i)_{\min})$ and $((p_{i+d})_{\min}, (q_{i+d})_{\min})$ coincide. Then

$$((p_{i+k})_{\min}, (q_{i+k})_{\min}) = ((p_{i+d+k})_{\min}, (q_{i+d+k})_{\min})$$

for every $k \geq 1$. In particular,

$$((p_{N_2})_{\min}, (q_{N_2})_{\min}) = ((p_{N_2+d})_{\min}, (q_{N_2+d})_{\min}).$$

It follows that

$$((p_{N_2})_{\min}, (q_{N_2})_{\min}) = ((p_{N_2+dn})_{\min}, (q_{N_2+dn})_{\min})$$

for every $n \geq 1$.

Thus, we can take $T := N_2$, $p := N_2!$. $\square$

Definition 7.6. Let τ be a reduced edge-path in G_r with a unique r -cancellation point y and suppose that y is non-deletable. The value

$$a := \sum_{i=1}^{\infty} L_r(\alpha_i) / \lambda^i, \quad (7.2)$$

where the paths α_i were defined in Definition 7.1, is called the *cancellation radius* for y in τ .

This terminology is justified by the following observation.

Observation 1. Let $\tau \equiv \bar{p}_0 \cdot q_0$ be a reduced edge-path in G_r , where the paths p_0, q_0 are r -legal and $y = \alpha(p_0) = \alpha(q_0)$ is a non-deletable r -cancellation point in τ . Let a be the cancellation radius of y in τ . Then, for each point $z \in H_r$ in τ , the following statements are equivalent:

- (1) The L_r -distance from z to y along the path τ is less than a .
- (2) There exists a natural number $N = N(z)$, such that the point $f^N(z)$ belongs to the interior of $I([f^N(p_0)], [f^N(q_0)])$. (So, the occurrence of $f^N(z)$ in $f^N(\tau)$ does not belong to the reduced path $[f^N(\tau)]$).

Observation 2. Let τ be a path and $y \in \tau$ be a vertex as in Definition 7.6. For $n \geq 0$, let y^n be the r -cancelation point in $[f^n(\tau)]$, i.e. y^n is the n -th successor of y . Let $a(n)$ be the cancelation radius of y^n in $[f^n(\tau)]$. Then, using (7.2), we have

$$a(n) = \sum_{i=1}^{\infty} L_r(\alpha_{i+n})/\lambda^i,$$

hence

$$\begin{aligned} \lambda^n a &= \lambda^n \left(\sum_{i=1}^n L_r(\alpha_i)/\lambda^i + \sum_{i=n+1}^{\infty} L_r(\alpha_i)/\lambda^i \right) = \\ &= L_r(f^{n-1}(\alpha_1)) + \cdots + L_r(f(\alpha_{n-1})) + L_r(\alpha_n) + a(n). \end{aligned} \quad (7.3)$$

Proposition 7.7. *Let τ be a reduced edge-path in G_r with a unique r -cancelation point y and suppose that y is non-deletable. Then the following statements hold.*

- 1) *We can algorithmically compute the r -cancelation radius a for y in τ .*
- 2) *$(p_0)_{\min}$ and $(q_0)_{\min}$ are the minimal initial edge-subpaths of p_0 and q_0 with last edges in H_r and L_r -length at least a .*

In particular, we can compute the pair $((p_0)_{\min}, (q_0)_{\min})$.

Proof. By Proposition 7.5, one can compute $T \geq 0$ and $p \geq 1$, such that for every $j \geq 1$ the paths α_{T+j} and α_{T+j+p} coincide. So,

$$a = \sum_{i=1}^T L_r(\alpha_i)/\lambda^i + \frac{\lambda^p}{\lambda^p - 1} \cdot \sum_{i=T+1}^{T+p} L_r(\alpha_i)/\lambda^i, \quad (7.4)$$

in particular, a is computable. The second statement follows from Definition 7.3 of $((p_0)_{\min}, (q_0)_{\min})$ and from the first statement of Lemma 7.4. $\square$

Notation. Let a, b, c, d be edge-paths in Γ . We write $(a, b) \subseteq_{\text{init}} (c, d)$ if a is an initial subpath of c and b is an initial subpath of d .

The proof of the following lemma is easy and we leave it for the reader.

Lemma 7.8. *Let τ be a reduced edge-path in G_r with a unique r -cancelation point y and suppose that y is non-deletable. Then, for every $i, k \geq 0$, we have*

$$((p_{i+k})_{\min}, (q_{i+k})_{\min}) \subseteq_{\text{init}} \Lambda([f^k((p_i)_{\min})], [f^k((q_i)_{\min})]). \quad (7.5)$$

Proposition 7.9. *Let τ be a reduced edge-path in G_r and suppose that y is a unique r -cancelation point in τ . Then we can algorithmically verify, whether y is deletable or not.*

Proof. Let T, p be the integer numbers as in Proposition 7.5. We compute $\tau_T = [f^T(\tau)]$ and denote $y_T := \alpha(p_T) = \alpha(q_T)$. First we check, whether y_T is an r -cancelation point in τ_T . If not, then y is deletable. Suppose that y_T is an

r -cancellation point in τ_T . We claim that y_T (and hence y) is non-deletable if and only if there exists a *witness* pair, i.e. a pair

$$(a, b) \underset{\text{init}}{\subseteq} (p_T, q_T)$$

such that the first and the last edges of a and b exist and lie in H_r , and

$$(a, b) \underset{\text{init}}{\subseteq} \Lambda([f^p(a)], [f^p(b)]).$$

Indeed, if y_T is non-deletable, then, by Lemma 7.8 and Proposition 7.5, and by the first statement of Lemma 7.4, we can set

$$(a, b) := ((p_T)_{\min}, (q_T)_{\min}).$$

The other direction is obvious. Moreover, one can decide algorithmically, whether a witness pair exists. $\square$

Proposition 7.10. *There exists a computable constant $S = S(f)$ such that for every reduced edge-path $\tau \subset G_r$ with a unique r -cancellation point y the following holds:*

if each of the paths $\tau, [f(\tau)], \dots, [f^S(\tau)]$ contains an r -cancellation point, then all the paths $\tau, [f(\tau)], \dots, [f^S(\tau)]$ contain r -cancellation points.

In particular, if $[f^S(\tau)]$ contains an r -cancellation point, then y is non-deletable.

Proof. The proof follows from the proof of Proposition 7.5; we can set $S = T + p$. $\square$

Lemma 7.11. *Let μ be a reduced edge-path in Γ . Then for each pair $(u, v) \in \Gamma^0 \times \Gamma^0$ with $f(u) = \alpha(\mu)$ and $f(v) = \omega(\mu)$, there exists a unique reduced edge-path σ in Γ with $\alpha(\sigma) = u$ and $\omega(\sigma) = v$ and with the property $[f(\sigma)] = \mu$. One can algorithmically find such σ .*

Proof. Let σ_1 be an arbitrary path from v to u . Then $\mu f(\sigma_1)$ is a closed path based at $\alpha(\mu)$. Since f is a homotopy equivalence, we can find a closed path τ based at u such that $[f(\tau)] = [\mu f(\sigma_1)]$. Then we can set $\sigma := [\tau \overline{\sigma_1}]$. The uniqueness of the desired σ follows from the fact that f is a homotopy equivalence. $\square$

Proposition 7.12. *There exists a computable constant $R = R(f)$ with the following property: Let τ be a reduced edge-path in G_r with a unique r -cancellation point y and suppose that y is non-deletable. Then the number of edges in $(p_0)_{\min}$ and in $(q_0)_{\min}$ is at most R .*

Proof. We write $(p_0)_{\min} = a_1 b_1 \dots a_s b_s a_{s+1}$, where all a_i are nontrivial edge-paths in H_r and all b_i are nontrivial edge-paths in G_{r-1} . By Lemma 7.4, the number of r -edges in $(p_0)_{\min}$ is bounded by n_{critical} . In particular, $s < n_{\text{critical}}$. So, it suffices to bound the number of edges in each b_i .

By Claim 3 in the proof of Proposition 7.5, there exists a computable natural number $k_2 = k_2(f)$ such that the segment b_s , and hence all segments b_i , can be moved into I_{k_2} in k_2 steps.

For $i = 1, \dots, s$, let $n(i)$ be the minimal number such that $f^{n(i)}$ moves b_i into $I_{n(i)}$; in particular $n(i) \leq k_2$. Then $[f^{n(i)-1}(b_i)]$ is present in $[f^{n(i)-1}(\tau)]$, but it is canceled out after applying f to $[f^{n(i)-1}(\tau)]$ and tightening. Hence, $l([f^{n(i)}(b_i)]) \leq C_*$, where C_* is the Cooper constant from Lemma 4.12.

By Lemma 7.11, the number of reduced edge-paths in Γ which can be mapped by a given homotopy equivalence followed by tightening to a given reduced edge-path (of l -length at most C_*) in Γ is at most $|\Gamma^0|^2$. Moreover, these paths can be algorithmically found. This gives a universal (depending only on f) bound on their l -lengths. So, the l -lengths of all b_i are bounded. $\square$

Definition 7.13. Let τ be a reduced edge-path in G_r with a unique r -cancellation point y and suppose that y is non-deletable. The r -cancellation area for y in τ is the subpath μ of τ which starts and ends on nontrivial r -segments (parts of r -edges) and can be presented as $\mu \equiv \mu_1\mu_2$ with $\omega(\mu_1) = \alpha(\mu_2) = y$, and $L_r(\mu_1) = L_r(\mu_2) = a$, where a is the cancellation radius of y in τ .

Corollary 7.14. *There are only finitely many r -cancellation radii and r -cancellation areas for f . These radii and areas can be computed.*

Proof. First we list all reduced edge-paths in G_r of length at most $2R$. Then we list those of them, which have a unique r -cancellation point and this point is non-deletable (see Proposition 7.9). For each of these paths we compute the corresponding r -cancellation radius (Proposition 7.7) and r -cancellation area. $\square$

8. STABLE r -PATHS AND THEIR A -DECOMPOSITIONS

Let Γ be a finite connected graph, $f : \Gamma \rightarrow \Gamma$ be a relative train track and let $\emptyset = G_0 \subset \dots \subset G_N = \Gamma$ be the maximal filtration in Γ . In this section, we fix a natural number $0 < r \leq N$ and assume that H_r is an exponential stratum. Let λ be the eigenvalue corresponding to H_r . We will analyze cancellations in f -images of paths in G_r with several r -cancellation points.

Definition 8.1. For any reduced edge-path $\tau \subset G_r$, let $P_r(\tau)$ be the number of r -cancellation points in τ . Clearly, $P_r([f^i(\tau)]) \geq P_r([f^{i+1}(\tau)])$ for any $i \geq 0$. We will say that τ is r -stable if $P_r([f^i(\tau)]) = P_r(\tau)$ for all $i \geq 1$. If τ is r -stable, we call the r -cancellation points of τ *non-deletable*. Clearly, there exists a nonnegative integer i_0 such that the path $[f^{i_0}(\tau)]$ is r -stable. We denote by $N_r(\tau)$ the number of r -cancellation points in $[f^{i_0}(\tau)]$.

We will show that i_0 and hence $N_r(\tau)$ are computable.

Proposition 8.2. *For any reduced edge-path $\tau \subset G_r$, one can compute i_0 such that the path $[f^{i_0}(\tau)]$ is r -stable. In particular, one can check, whether τ is r -stable or not.*

Proof. Let $y_1, \dots, y_k$ be all r -cancellation points in τ . Let y_0 and y_{k+1} be the initial and the terminal vertices of τ , respectively. Let τ_i be the subpath of τ from y_i to y_{i+1} , $i = 0, \dots, k$.

First we note that y_i is the unique r -cancellation point in the edge-path $\tau_{i-1}\tau_i$. For each $1 \leq i \leq k$, we can check in S steps by Proposition 7.10, whether the point y_i in $\tau_{i-1}\tau_i$ is deletable or not. If it is deletable for some i , then $P_r([f^S(\tau)]) < P_r(\tau)$ and we can proceed by induction restarting from $[f^S(\tau)]$. Suppose that y_i is non-deletable r -cancellation point in $\tau_{i-1}\tau_i$ for each $1 \leq i \leq k$. By Proposition 7.7, we can compute the r -cancellation radius of y_i in $\tau_{i-1}\tau_i$. Let a_i be this radius.

If $L_r(\tau_i) \geq a_i + a_{i+1}$ for $i = 1, \dots, k-1$, then the path τ is stable and the points $y_1, \dots, y_k$ are non-deletable.

Suppose that for some i we have $L_r(\tau_i) < a_i + a_{i+1}$. Let τ'_i, τ''_i be two subpaths of τ_i such that $\tau_i \equiv \tau'_i\tau''_i$, $L_r(\tau'_i) < a_i$, and $L_r(\tau''_i) < a_{i+1}$. Then there exists K' such that the path $[f^{K'}(\tau_{i-1}\tau'_i)]$ is r -legal and there exists K'' such that the path $[f^{K''}(\tau''_i\tau_{i+1})]$ is r -legal. We can find $K = \max\{K', K''\}$ by brute force (although there is a way to estimate K in terms of f). For this K we have $N_r([f^K(\tau)]) < N_r(\tau)$ and again we can proceed by induction. $\square$

For convenience, we reformulate a part of Definition 8.1.

Definition 8.3. Let τ be a reduced edge-path in G_r and let $y_1, \dots, y_k$ be all r -cancellation points in τ . We say that the r -cancellation points in τ are *non-deletable* if the number of r -cancellation points in $[f^i(\tau)]$ is equal to k for every $i \geq 0$.

Proposition 8.4. (*Criterion of r -stability*) *Let τ be a reduced edge-path in G_r and let $y_1, \dots, y_k$ be all r -cancellation points in τ . Let y_0 and y_{k+1} be the initial and the terminal vertices of τ , respectively. Let τ_i be the subpath of τ from y_i to y_{i+1} . Then all cancellation points $y_1, \dots, y_k$ are non-deletable in τ if and only if each of y_i is non-deletable in $\tau_{i-1}\tau_i$ for $i = 1, \dots, k$, where a_i is the cancellation radius for y_i in $\tau_{i-1}\tau_i$.*

The proof follows from the proof of Proposition 8.2.

Definition 8.5. With notations in Proposition 8.4, suppose that the r -cancellation points $y_1, \dots, y_k$ in τ are non-deletable in τ . Let A_i be the r -cancellation area of y_i in $\tau_{i-1}\tau_i$ for $i = 1, \dots, k$. Then $A_1, \dots, A_k$ are called the *r -cancellation areas* of $y_1, \dots, y_k$ in τ .

Definition 8.6. Let τ be a reduced edge-path in G_r whose r -cancellation points are non-deletable. Let $y_1, \dots, y_k$ be these cancellation points and $A_1, \dots, A_k$ be the corresponding r -cancellation areas in τ . Then we can write $\tau \equiv b_0 \cdot A_1 \cdot b_1 \cdot \dots \cdot A_k \cdot b_k$, where $b_0, \dots, b_k$ are some r -legal paths in G_r . We call such expression the *A-decomposition* of τ .

Remark 8.7. Suppose that τ has the *A-decomposition*

$$\tau \equiv b_0 \cdot A_1 \cdot b_1 \cdot \dots \cdot A_k \cdot b_k.$$

Then, for every $i \geq 1$, the path $[f^i(\tau)]$ has the *A-decomposition*

$$[f^i(\tau)] \equiv b_0^i \cdot A_1^i \cdot b_1^i \cdot \dots \cdot A_k^i \cdot b_k^i, \quad (8.1)$$

where $b_j^i \equiv [f^i(b_j)]$ and $A_j^i \equiv [f^i(A_j)]$ for all possible j .

9. SUBDIVIDED RELATIVE TRAIN TRACK

It is technically convenient to start the proof of the main theorem in the situation where the following condition is satisfied:

(RTT-iv) There is a computable natural number $P = P(f)$ such that for each exponential strata H_r and each r -cancellation area A of f , the r -cancellation area $[f^P(A)]$ is an edge-path.

We will show that this condition is satisfied for a map $f' : \Gamma' \rightarrow \Gamma'$ obtained from $f : \Gamma \rightarrow \Gamma$ by using subdivisions of edges of Γ at certain points. Moreover, f' with respect to a natural filtration on Γ' will satisfy the properties (RTT-i) – (RTT-iii) of a relative train track.

Definition 9.1. Let H_r be an exponential strata for the relative train track $f : \Gamma \rightarrow \Gamma$ and let $A_1, \dots, A_s$ be all r -cancellation areas of f . An endpoint v of A_i is called an *r-subdivision point* if v is not a vertex (hence v lies in the interior of some edge $e(v)$ from H_r) and $f^n(v)$ is not a vertex for all natural n .

Proposition 9.2. The following claims are valid:

- 1) The map f acts on the set of the r -cancellation areas by sending A_i to $[f(A_i)]$, in particular f acts on the set of endpoints of these areas.
- 2) One can find the set of r -subdivision points.
- 3) The set of r -subdivision points is f -invariant.
- 4) If v is an endpoint of an r -cancellation area and v is not an r -subdivision point, then $f^{(2s-1)!}(v)$ is a vertex of Γ . Here s is the number of r -cancellation areas.

Proof. Claims 1) and 3) are obvious. We prove the claim 4). By Corollary 7.14, we can find all r -cancellation areas $A_1, \dots, A_s$ and hence all their endpoints $v_1, \dots, v_{2s}$. Let v be one of these endpoints. Since f acts on the set $\{v_1, \dots, v_{2s}\}$, v is not an r -subdivision point if and only if $f^i(v)$ is a vertex for some $0 \leq i \leq 2s - 1$. This completes the proof of the claim 4). The claim 2) follows from the claim 4). $\square$

Definition 9.3. Let Γ' be the graph obtained from Γ by subdivision at all r -subdivision points for all exponential strata H_r . By Claim 2), we can construct this graph efficiently. By Claim 3), the map $f : \Gamma \rightarrow \Gamma$, induces a map $f' : \Gamma' \rightarrow \Gamma'$. We also define a natural filtration $\emptyset = G'_0 \subset \dots \subset G'_N = \Gamma'$ by saying that G'_i consists of all edges resulting from G_i by the subdivision. It is possible that this filtration is not maximal. A stratum H'_i is called exponential, polynomial, or zero if the strata H_i is exponential, polynomial, or zero, respectively. For any exponential stratum H'_r we have the induced pseudo-metric L'_r on G'_r . It is possible that $L_r(e) = 0$ for some edge $e \in H'_r$. However, this will not destroy the further proof. Note that the r -cancellation areas of f and f' can be naturally identified.

One can show that the homotopy equivalence $f' : \Gamma' \rightarrow \Gamma'$ satisfies the conditions (RTT-i) – (RTT-iii) in the definition of a relative train track. Therefore f' is called the *subdivided relative train track* associated with f .

The advantage of f' is that also the property (RTT-iv) for f' is satisfied.

Each edge e in Γ can be considered in Γ' as an edge-path. Let $n(e)$ be the length of this path. We set

$$n_f := \max\{n(e) : e \in \Gamma^1\}.$$

Proposition 9.4. *Let τ be a reduced f' -path in Γ' of length at least $n_f + n_f \|f'\|$. If τ lies in G'_r , then there exists a natural $k \leq n_f$ such that $\widehat{f'}^k(\tau) \subset G'_{r-1}$ or $L_r(\widehat{f'}^k(\tau)) > 0$.*

Proof. Write $\tau \equiv e_1 \dots e_{n_f} \dots e_s$, where e_i are edges of Γ' . By definition of n_f , there exists $k \leq n_f$ such that $\omega(e_k)$ is a vertex of Γ . Then $p := \widehat{f'}^k(\tau)$ is a path in G'_r with endpoints in Γ^0 . Then either p lies in G'_{r-1} or p contains an edge of Γ from H_r . In the latter case we have $L_r(p) > 0$. $\square$

Agreement. From now on we will work with f' and never with the old f . So, we will simplify notation and skip dashes by writing f, Γ, G_i, H_i instead of f', Γ', G'_i, H'_i , respectively.

Below we collect the properties of the new f which we will use later.

Proposition 9.5. *The subdivided relative train track $f : \Gamma \rightarrow \Gamma$ satisfies the following properties:*

- 1) *This f represents the same automorphisms of F as the original one.*
- 2) *This f satisfies the condition (Pol) as the original one.*
- 3) *For each exponential stratum H_r , there exists only finitely many r -cancellation areas. These areas can be computed.*
- 4) *Each r -cancellation area has an initial and a terminal subpaths which lie in H_r and have nonzero L_r -lengths.*
- 5) *If A is an r -cancellation area in a reduced path $\tau \subset G_r$, then $[f(A)]$ is an r -cancellation area in $[f(\tau)]$.*
- 6) *One can compute a natural number $R_\star = R_\star(f)$ such that each r -cancellation area of f is divided by the corresponding non-deletable r -cancellation point into two parts such that each part is contained in an edge-path of l -length at most $R_\star$.*
- 7) *One can compute a natural number $P = P(f)$ such that for every exponential stratum H_r and every r -cancellation area A , the r -cancellation area $[f^P(A)]$ is an edge-path.*
- 8) *One can compute a natural number $K = K(f)$ such that the following holds: Let H_r be an exponential stratum and τ be a reduced f -path in G_r of l -length at least K . Then there exists a natural $k \leq K$ such that either $\widehat{f}^k(\tau) \subset G_{r-1}$, or $L_r(\widehat{f}^k(\tau)) > 0$.*

10. r -PERFECT AND A -PERFECT PATHS

Definition 10.1. Let H_r be an exponential stratum. An edge-path $\tau \subset G_r$ is called *r -perfect* if the following conditions are satisfied:

- (i) τ is a reduced f -path and its first edge belongs to H_r ,
- (ii) τ is r -legal,
- (iii) $[\tau f(\tau)] \equiv \tau \cdot [f(\tau)]$ and the turn of this path at the point between τ and $[f(\tau)]$ is legal.

Note that these conditions imply that $[\tau f(\tau)]$ is r -legal. In the following proposition we formulate some important properties of r -perfect paths; they can be proved directly from the above definition.

Proposition 10.2. *Let τ be an r -perfect path. Then the following statements hold:*

- (1) *For every $i \geq 0$, the path $[f^i(\tau)]$ is r -perfect.*
- (2) *For every $i \geq 0$, the vertex $[f^i(\tau)]$ of D_f lies in the τ -subgraph. Moreover, $[f^i(\tau)] \equiv \widehat{f}^{m_i}(\tau)$ for some computable m_i satisfying $m_i < m_{i+1}$.*
- (3) *For every $i \geq 0$, the path $\widehat{f}^i(\tau) \subset G_r$ is r -legal, contains edges from H_r , and $L_r(\widehat{f}^{i+1}(\tau)) \geq L_r(\widehat{f}^i(\tau))$.*

Definition 10.3. Let H_r be an exponential stratum. A reduced f -path $\tau \subset G_r$ containing edges from H_r is called *A-perfect* if

- (i) all r -cancelation points in τ are non-deletable,
the corresponding r -cancelation areas are edge-paths,
- (ii) the A -decomposition of τ starts on an A -area, i.e. it has the form
 $\tau \equiv A_1 b_1 \dots A_k b_k$,
- (iii) $[\tau f(\tau)] \equiv \tau \cdot [f(\tau)]$ and the turn at the point between τ and $[f(\tau)]$ is legal.

The following proposition can be proved straightforward and we leave it for the reader.

Proposition 10.4. Let H_r be an exponential stratum and let τ be an A -perfect path in G_r with the A -decomposition $\tau \equiv A_1 b_1 \dots A_k b_k$.

For $1 \leq j \leq k$, we set $\tau_{0,j} \equiv [A_j b_j \dots A_k b_k f(A_1 b_1 \dots A_{j-1} b_{j-1})]$ and for $i \geq 1$ we set $\tau_{i,j} \equiv [f^i(\tau_{0,j})]$. Then the following statements are satisfied:

- (1) For any $1 \leq j \leq k$ and $i \geq 0$ the path $\tau_{i,j}$ is A -perfect.
- (2) For any $1 \leq j \leq k$ and $i \geq 0$ the vertex $\tau_{i,j}$ of D_f lies in the τ -subgraph.
Moreover $\tau_{i,j} \equiv \widehat{f}^{m_{i,j}}(\tau)$ for some computable $m_{i,j}$ satisfying $m_{i,j} < m_{i,j+1}$
and $m_{i,k} < m_{i+1,1}$.
- (3) Every vertex σ from the τ -subgraph such that the path σ in G_r is A -perfect coincides with $\tau_{i,j}$ for some i, j .
- (4) For every vertex σ in the τ -subgraph, at least one of the paths $\sigma, \widehat{f}(\sigma), \dots, \widehat{f}^{l(\sigma)}(\sigma)$ coincides with $\tau_{i,j}$ for some i, j .

11. SOME TECHNICAL PROPOSITIONS

Proposition 11.1. Let H_r be an exponential stratum. Let μ be a reduced r -legal f -path with endpoints in H_r and with the first edge in G_{r-1} , and the last edge in H_r . Then the μ -subgraph of D_f contains a vertex which is an r -perfect path in G_r . This vertex can be algorithmically found.

Proof. We write $\mu \equiv \sigma\tau$, where σ is a nontrivial path in G_{r-1} and τ is a nontrivial path in G_r with the first and the last edges from H_r . We prove that $\mu' := \widehat{f}^{l(\sigma)}(\mu)$ is r -perfect and lies in the μ -subgraph.

Clearly, $\mu' \equiv \tau[f(\sigma)]$, μ' is r -legal and lies in the μ -subgraph. It remains to prove that the turn between μ' and $[f(\mu')]$ in $\mu' \cdot [f(\mu')]$ is legal.




Figure 10.

By (RTT-ii), $[f(\sigma)]$ is a non-degenerate path in G_{r-1} . So the last edge of μ' lies in G_{r-1} . On the other hand, the first edge of $[f(\mu')] \equiv [f(\tau)][f^2(\sigma)]$ lies in H_r . Hence, the turn under consideration is mixed and hence legal. $\square$

Proposition 11.2. *Let H_r be an exponential stratum and let $\mu \subset G_r$ be a reduced f -path such that the paths μ and $[\mu f(\mu)]$ are r -legal. Then there exists a vertex μ' of the graph D_f which is contained in the μ -subgraph and for which at least one of the following statements is satisfied:*

- (1) *the path μ' is contained in the subgraph G_{r-1} ;*
- (2) *the path μ' is r -perfect;*
- (3) *the μ' -subgraph is finite.*

Moreover, there exists an algorithm, which constructs such a vertex μ' and indicates one of the statements (1)-(3) which is satisfied for μ' .

Proof. We assume that μ contains an edge or a partial edge from H_r , otherwise we can set $\mu' = \mu$ and (1) is satisfied.

Write $\mu \equiv b_1 \cdot b_2 \cdot b_3$, where the edge-paths (possibly trivial) b_1, b_3 lie in G_{r-1} , the path b_2 begins and ends on edges from H_r . We claim that the path $\tilde{\mu} := b_2 \cdot [b_3 f(b_1)]$ satisfies the following properties:

- i) The path $\tilde{\mu}$ is a reduced f -path, which begins on a (partial) edge from H_r ; it has the same number of edges and partial edges from H_r as μ .
- ii) The vertex $\tilde{\mu}$ of D_f lies in the μ -subgraph.
- iii) The paths $\tilde{\mu}$ and $[\tilde{\mu} f(\tilde{\mu})]$ are r -legal.

Indeed, i) is obvious; ii) is fulfilled, since $\tilde{\mu} \equiv \hat{f}^{l(b_1)}(\mu)$.

The claim iii) follows from the formulas

$$\tilde{\mu} \equiv [[\bar{b}_1]\mu[f(b_1)]], \quad [\tilde{\mu}f(\tilde{\mu})] \equiv [[\bar{b}_1][\mu f(\mu)][f^2(b_1)]],$$

from the assumption that μ and $[\mu f(\mu)]$ are r -legal, and from the observation that the product of an r -legal path and a path from G_{r-1} remains r -legal.

Thus, replacing μ by $\tilde{\mu}$, we may additionally assume that μ begins on a (partial) edge E from H_r . Since μ is r -legal, $f(E)$ is an initial subpath of $[f(\mu)]$. Let τ be the initial subpath of μ such that $\mu \equiv \tau \cdot \overline{I(\bar{\mu}, [f(\mu)])}$. We consider the following four cases (see Figure 11).

Case (1) The path τ is non-degenerate (hence, it contains E) and $f(E)$ is an initial subpath of $I(\bar{\mu}, [f(\mu)])$.

Case (2) The path τ is non-degenerate (hence, it contains E) and $I(\bar{\mu}, [f(\mu)])$ is a proper initial subpath of $f(E)$.

Case (3) The path τ is degenerate (hence, $\mu = \overline{I(\bar{\mu}, [f(\mu)])}$) and $f(E)$ is a proper initial subpath of $I(\bar{\mu}, [f(\mu)])$.

Case (4) The path τ is degenerate (hence, $\mu = \overline{I(\bar{\mu}, [f(\mu)])}$) and $I(\bar{\mu}, [f(\mu)])$ is an initial subpath of $f(E)$.




Figure 11.

In (1) and (2), the path μ is depicted by the large left corner, and in (3), (4), it is depicted by a vertical line. The path $[f(\mu)]$ is depicted by the large right corner.

Consider Cases (1) and (3).

Let $\mu_1 := [\overline{E}\mu f(E)]$. Clearly, μ_1 lies in the μ -subgraph. Moreover, the paths μ_1 and $[\mu_1 f(\mu_1)]$ are r -legal. Indeed, in these subcases μ_1 is a subpath of the r -legal path μ ; the path $[\mu_1 f(\mu_1)]$ is a subpath of the r -legal path μ or $[\mu f(\mu)]$ in Case (1), and it is a subpath of the r -legal path $[f(\mu)]$ in Case (3). Since the number of r -edges in μ_1 is less than in μ , we can proceed by induction and find the desired μ' in the μ_1 -subgraph.

Consider Case (2). Let E_1 be the second edge of $\mu[f(\mu)]$. In Figure 12, we distinguish the cases where τ contains at least two edges (picture (1)) and where τ contains exactly one edge (picture (2)).

We set $\mu_1 \equiv [\overline{E}\mu f(E)]$. The path μ_1 is r -legal as a subpath of $[\mu f(\mu)]$, it begins on E_1 and ends on the last edge of $f(E)$, and it lies in the μ -subgraph.

Suppose that E_1 is an edge from H_r . Then $[f(\mu_1)]$ begins on the first edge of $f(E_1)$. Therefore the turn between μ_1 and $[f(\mu_1)]$ coincides with the turn between $f(E)$ and $f(E_1)$ which is legal (since the turn between E and E_1 is a turn of the r -legal path $[\mu f(\mu)]$). Hence, μ_1 is r -perfect and we have the statement (2) for $\mu' \equiv \mu_1$.

Now suppose that E_1 lies in G_{r-1} . By Proposition 11.1 applied to μ_1 , we can construct μ' satisfying the statement (2).




Figure 12.

Consider Case (4). Let $\mu \equiv E \cdot \sigma$, then $f(E) \equiv \bar{\sigma} \cdot \overline{E} \cdot \delta$ for some path δ . This implies $\widehat{f}(\mu) \equiv \overline{E} \cdot \delta$ and $\widehat{f}^2(\mu) \equiv \mu$. Hence, the μ -subgraph is finite and for $\mu' \equiv \mu$ we have the statement (3). $\square$

Recall that $C_\star$ is the Cooper constant from Lemma 4.12 and $R_\star$ is the constant from Proposition 9.5.

Proposition 11.3. *Let H_r be an exponential stratum. Let $\sigma \subset G_r$ be a reduced f -path which is r -legal. Suppose that $[\sigma f(\sigma)]$ contains a non-deletable r -cancellation point and the r -cancellation area in $[\sigma f(\sigma)]$ is an edge-path. Then the σ -subgraph contains a vertex σ' for which at least one of the following statements is satisfied:*

- (1) $l(\sigma') \leq C_\star + 2R_\star$;
- (2) the path σ' is A -perfect.

Moreover, there exists $k \leq l(\sigma)$ such that $\sigma' \equiv \widehat{f}^k(\sigma)$ satisfies (1) or (2). In particular, such σ' can be algorithmically found.

Proof. Let z^i be the unique non-deletable r -cancellation point of the path $[f^i(\sigma)f^{i+1}(\sigma)]$ and let $A(z^i)$ be the r -cancellation area of z^i in this path, $i = 0, 1$. We have $A(z^1) \equiv [f(A(z^0))]$. So, if a and b are the initial and the terminal vertices of $A(z^0)$, then $f(a)$ and $f(b)$ are the initial and the terminal vertices of $A(z^1)$.

For two vertices u, v in $[f(\sigma)]$, we write $u < v$ if u and v are the initial and the terminal vertices of some nontrivial subpath of $[f(\sigma)]$. If $z^1 \leq z^0$, we can find σ' satisfying (1). Let $z^0 < z^1$.

We write $\sigma \equiv p \cdot q$, where p is the initial segment of σ until the vertex a . Observe that, by Cooper's lemma, if a point t moves along p from $\alpha(\sigma)$ to a , then $f(t)$ moves in the $C_\star$ -neighborhood of $[f(p)]$. This implies the following claim.

Claim. Let $p \equiv p_1 p_2$, where p_1, p_2 are edge-paths. If $p_2 q$ completely cancels in $p_2 q \cdot [f(p_1)]$, then $l([p_2 q \cdot [f(p_1)]]) \leq C_\star$.




Figure 13.

Suppose that there is a decomposition $p \equiv p_1 p_2$ such that $p_2 q$ completely cancels in $p_2 q \cdot [f(p_1)]$ and that p_1 is the shortest initial segment of p with this property.

We set $\sigma' := \hat{f}^{l(p_1)}(\sigma)$. Then $\sigma' \equiv [p_2 q \cdot [f(p_1)]]$, so $l(\sigma') \leq C_\star$ by the above claim. Obviously, σ' lies in the σ -subgraph. Thus, in this case we have the statement (1).

Now suppose that cancellations in $p_2 q \cdot [f(p_1)]$ don't touch the first edge of p_2 for every initial segment p_1 of p . We set $\sigma' := \hat{f}^{l(p)}(\sigma)$. Then $\sigma' \equiv q \cdot [f(p)]$. Consider two cases.

Case 1. Suppose that $b < f(a)$ (see Figure 13).

Then σ' is the subpath of $[\sigma f(\sigma)]$ from a to $f(a)$. We prove that σ' is A -perfect. Then the statement (2) will be fulfilled. First note that σ' has the A -decomposition $\sigma' \equiv A(z^0) \cdot \ell$, where ℓ is the subpath of $[f(\sigma)]$ from b to $f(a)$. We have $[f(\sigma')] \equiv A(z^1) \cdot [f(\ell)]$. By Lemma 7.4, the first edge of the r -cancellation area $A(z^1)$ lies in H_r . Then, in $\sigma' \cdot [f(\sigma')]$, the turn at the point between σ' and $[f(\sigma')]$ is either mixed in (G_{r-1}, G_r) or is an r -turn. In particular, this turn is legal, since it belongs to the r -legal path $[f(\sigma)]$.

Case 2. Suppose that $f(a) \leq b$.

We prove that in this case we have the statement (1). Consider two subcases.

Subcase 1. Suppose that $f(a)$ lies on the subpath of $[f(\sigma)]$ from z^0 to b . Then $l(\sigma') = l(a, f(a)) \leq l(a, b) \leq 2R_\star$.

Subcase 2. Suppose that $f(a)$ lies on the subpath of $[f(\sigma)]$ from $\alpha([f(\sigma)])$ to z^0 . Then $l(\sigma') = l(a, f(a)) \leq l(a, z^0) + l(f(a), z^0) \leq l(a, z^0) + l(f(a), z^1) \leq 2R_\star$. $\square$

12. INFINITE μ -SUBGRAPHS AND f -ORBITS OF SOME VERTICES OF D_f

In this section we use the following notation. Let M be a subtree of a tree $\widetilde{M}$. Every vertex $v \in \widetilde{M}$ can be naturally projected to M ; we denote this projection by $\text{Pr}_M(v)$. The edge-distance between vertices $u, w \in \widetilde{M}$ is denoted by $\text{dist}(u, w)$.

Proposition 12.1. *Let μ be a vertex of D_f such that the following conditions are satisfied:*

- i) *the μ -subgraph (we denote it by M) is infinite,*
- ii) *$l(\tau) > C_\star$ for every vertex τ in M ,*
- iii) *μ divides the graph D_f into at least two components, and the closure of the component containing M is a tree. We denote this closure by $\widetilde{M}$.*

Then the f -paths $[f^i(\mu)]$, $i \geq 1$, considered as vertexes of D_f , lie in $\widetilde{M}$ and with notation $\tau_i = \text{Pr}_M([f^i(\mu)])$ we have

$$\text{dist}(\mu, \tau_i) < \text{dist}(\mu, \tau_{i+1}).$$

Proof. Let $\mu \equiv E_1 E_2 \dots E_k$ and let p_μ be the path in D_f which starts at the vertex μ and has labels of edges $E_1, \dots, E_k$. The consecutive vertices of p_μ are

$$\begin{aligned} & E_1 E_2 \dots E_k, \\ & [E_2 E_3 \dots E_k f(E_1)], \\ & [E_3 E_4 \dots E_k f(E_1) f(E_2)], \\ & \vdots \\ & [f(E_1) f(E_2) \dots f(E_k)]. \end{aligned}$$




Figure 14.

Since E_1 is the first edge of μ , the first edge of p_μ lies in the μ -subgraph. (Note that E_2 is not necessarily the first edge of $[\overline{E_1}\mu f(E_1)]$, so the second edge of p_μ possibly does not lie in the μ -subgraph.)

Since $E_1 E_2 \dots E_k$ is reduced and $\widetilde{M}$ is a tree, the path p_μ is without backtracking and hence its terminal vertex $[f(\mu)]$ lies in $\widetilde{M}$. By analogy, the path p_{τ_1} starts at τ_1 and ends at $[f(\tau_1)]$, it has label τ_1 , and its first edge lies in the τ_1 -subgraph, which is a part of M . By ii), we have

$$\text{dist}(\tau_1, [f(\tau_1)]) = l(\tau_1) > C_\star \quad (12.1)$$

Note that τ_1 lies on the reduced path p_μ connecting μ and $[f(\mu)]$. By Cooper's lemma, $[f(\tau_1)]$ is at distance at most $C_\star$ from some reduced path connecting $[f(\mu)]$ and $[f^2(\mu)]$. From this and (12.1) we have

$$\text{dist}(\mu, \tau_1) < \text{dist}(\mu, \tau_2),$$

see Figure 14. Note that τ_2 lies on a reduced path connecting $[f(\mu)]$ and $[f^2(\mu)]$. Continuing as above, we complete the proof. $\square$

Proposition 12.2. *Let μ be a vertex of D_f such that the following conditions are satisfied:*

- i) *the μ -subgraph (we denote it by M) is infinite,*
- ii) *$l(\tau) > C_\star$ for every vertex τ in M ,*
- iii) *μ divides the graph D_f into at least two components, and the closure of the component containing M is a tree. We denote this closure by $\widetilde{M}$.*

Let σ be an arbitrary vertex of $\widetilde{M}$. Then at least one of the vertices $[f(\sigma)]$, $[f^2(\sigma)]$ belongs to $\widetilde{M}$.

Proof. We denote $\tau_1 = \text{Pr}_M(\sigma)$. Suppose that $[f(\sigma)]$ does not belong to $\widetilde{M}$. We define the path p_σ in D_f as in Proposition 12.1. Then the path p_σ is reduced, has the label σ and it connects σ and $[f(\sigma)]$.




Figure 15.

Since σ belongs to $\widetilde{M}$ and $[f(\sigma)]$ not, the vertex τ_1 belongs to p_σ . Using Cooper's lemma, one can deduce that $[f(\tau_1)]$ is at distance at most $C_\star$ from some reduced path connecting $[f(\sigma)]$ and $[f^2(\sigma)]$. The proof can be completed as in Proposition 12.1, compare Figures 15 and 14. $\square$

13. GOOD VERTICES IN INFINITE μ -GRAPHS

Until the end of this section we will assume that H_r is an exponential stratum. Recall that a reduced edge-path $\tau \subset G_r$ is called r -stable if all r -cancellation points in τ are non-deletable (see Definitions 8.1 and 8.3).

By Proposition 8.2, for any reduced edge-path $\mu \subset G_r$, one can compute a natural number i_0 such that $[f^{i_0}(\mu)]$ is r -stable. We denote by $N_r(\mu)$ the number of r -cancellation points in $[f^{i_0}(\mu)]$. Note that $N_r(\mu) = N_r([f^i(\mu)])$ for every $i \geq 0$.

Definition 13.1. A reduced f -path $\tau \subset G_r$ is called r -superstable if all r -cancellation points in τ and in $[\tau f(\tau)]$ are non-deletable and all r -cancellation areas in these paths are edge-paths.

Note that if τ is r -superstable, then $[f^i(\tau)]$ is r -superstable for all $i \geq 0$.

Lemma 13.2. For any reduced f -path $\tau \subset G_r$, one can algorithmically compute a natural number S such that the path $[f^S(\tau)]$ is r -superstable.

Proof. By Propositions 8.2 and by the statement 7) of Proposition 9.5, we can compute a number S_1 such that all r -cancellation points in $[f^{S_1}(\tau)]$ are non-deletable and all r -cancellation areas in τ are edge-paths. Also, we can compute a

number S_2 such that all r -cancelation points in $[f^{S_2}([\tau f(\tau)])]$ are non-deletable and all r -cancelation areas in $[f^{S_2}([\tau f(\tau)])]$ are edge-paths. We set $S = \max\{S_1, S_2\}$. Then the path $[f^S(\tau)]$ is superstable. $\square$

Proposition 13.3. *Suppose that μ is a reduced f -path in G_r and suppose that the following conditions are satisfied:*

- i) *the μ -subgraph (we denote it by M) is infinite,*
- ii) *μ divides D_f into at least two components, and the closure of the component containing M is a tree. We denote this closure by $\widetilde{M}$.*
- iii) *$\widetilde{M}$ does not contain dead and repelling vertices, and vertices τ with $l(\tau) \leq 2(R_\star + C_\star)$.*

Then we can algorithmically find a vertex μ' in the μ -subgraph with one of the following properties:

- 1) *the f -path μ' lies in G_{r-1} ,*
- 2) *μ' is r -perfect,*
- 3) *μ' is A -perfect.*

Proof. First observe, if μ'' is a vertex in $\widetilde{M}$, then flowing along the μ'' -subgraph, we meet the μ -subgraph. In particular, these subgraphs (rays) coincide up to finite initial segments. This follows from the assumption that $\widetilde{M}$ does not contain dead and repelling vertices.

Therefore, if we will find $\mu'' \in \widetilde{M}$ with one of the properties 1)-3), then we will be able to find the desired μ' just by flowing along the μ'' -subgraph until the μ -subgraph and then by flowing along the μ -subgraph (see the statements (1) and (2) in Propositions 10.2 and 10.4).

(A) A plan for construction of $\mu'' \in \widetilde{M}$ with one of the properties 1)-3).

Step 1. First we find an r -superstable vertex μ_0 in $\widetilde{M}$. We can take, for instance, $\mu_0 := [f^S(\mu)]$, where S is the number from Lemma 13.2. By Proposition 12.1, μ_0 lies in $\widetilde{M}$ and obviously $N_r(\mu_0) = N_r(\mu)$. We go to Step 2 or 3 depending on whether $N_r(\mu_0)$ is zero or not.

Step 2. Suppose that $N_r(\mu_0) = 0$.

If $N_r([\mu_0 f(\mu_0)]) = 0$, we can find the desired μ'' by Proposition 11.2 (as described above, the μ'' -subgraph almost coincides with the μ -subgraph, so it cannot be finite).

If $N_r([\mu_0 f(\mu_0)]) = 1$, we can apply Proposition 11.3 to $\sigma := \mu_0$ and find μ'' in the μ_0 -subgraph such that one of the following cases holds:

- (1) $l(\mu'') \leq C_\star + 2R_\star$.
- (2) The path μ'' is A -perfect.

Case (1) is impossible by iii). In Case (2) we are done.

Step 3. Suppose that $N_r(\mu_0) \geq 1$. Then we will find in (B) an r -superstable vertex

$\mu_1 \in \widetilde{M}$ for which one of the following is satisfied:

- (1) $N_r(\mu_1) < N_r(\mu_0)$.
- (2) μ_1 is A -perfect.

In Case (1) we replace μ_0 by μ_1 and proceed by Step 2 or 3 again.
In Case (2) we are done.

Remark. Proposition 12.2 and Lemma 13.2 imply the following claim:

For every vertex τ in $\widetilde{M}$, one can find a natural number S such that $\tau' := [f^S(\tau)]$ lies in $\widetilde{M}$ and is r -superstable. Clearly, $N_r(\tau') = N_r(\tau)$. Thus, in the following we will not take care on r -superstability.

(B) A realization of the plan in Case $N_r(\mu_0) \geq 1$.

Let $y_1, \dots, y_k$ ($k \geq 1$) be all r -cancellation points in μ_0 (they are non-deletable, since μ_0 is r -stable) and let $y_1^i, \dots, y_k^i$ be their i -successors in $[f^i(\mu_0)]$. The terminal point of $I_i = I(\overline{[f^i(\mu_0)]}, [f^{i+1}(\mu_0)])$ is denoted by z^i . For two vertices $v, u \in \mu_0$, we denote by $l_{\mu_0}(v, u)$ the length of the subpath of μ_0 connecting v and u .

Case 1. Suppose that y_k lies in I_0 .

The proof is illustrated by Figure 16, where we distinguish two cases:

- (a) y_k lies in the interior of I_0 . In this case $y_k = y_1^1$.
- (b) y_k coincides with the terminal vertex of I_0 .

The point y_1 divides μ_0 into two subpaths which we denote by p and q , so $\mu_0 \equiv pq$.

Claim. For every decomposition $p \equiv p_1 p_2$, where p_1 is a proper initial edge-subpath of p , cancellations in the product $p_2 q \cdot [f(p_1)]$ don't touch the first edge of p_2 .

To prove this claim, it suffices to note that $[f(p_1)]$ is r -legal, but $p_2 q$ is not. The first follows from the fact that p and so p_1 are r -legal, the second from the fact that the r -cancellation point y_k belongs to $p_2 q$.

Now we set $\mu_1 \equiv \widehat{f}^{l(p)}(\mu_0)$. Clearly, μ_1 lies in the μ_0 -subgraph and so in $\widetilde{M}$. We assert that $N_r(\mu_1) < N_r(\mu_0)$. To prove this, we describe the path μ_1 precisely. First, by the above claim we have $\mu_1 \equiv [q f(p)]$. Second, we observe that $\mu_1 \equiv \mu_{y_1 y_k} \cdot \ell$, where $\mu_{y_1 y_k}$ is the initial subpath of q from y_1 to y_k and ℓ is the final subpath of $[f(p)]$ from y_k to $f(y_1)$. Since ℓ is r -legal, the r -cancellation points of μ_1 are contained in the set $\{y_1, \dots, y_k\} \setminus \{y_1\}$. So, $N_r(\mu_1) < N_r(\mu_0)$.




Figure 16.

The path μ_1 is depicted by the bold line. The point y_1 on the left picture can lie in I_0 . The point y_1^1 on the right picture can coincide with y_k .

Case 2. Suppose that y_k does not lie in I_0 , but y_1^1 lies in I_0 . In this case $y_1^1 = z^0$, where z^0 is the terminal vertex of I_0 .

Case 2.1. Suppose that $l_{\mu_0}(y_1, y_1^1) > R_\star + C_\star$. The proof in this case is illustrated by Figure 17.

Let u be the terminal vertex of the r -cancellation area $A(y_1)$. The point u divides μ into two subpaths which we denote by p and q , so $\mu_0 \equiv pq$.

By definition of $R_\star$, we have $l_{\mu_0}(y_1, u) \leq R_\star$, hence $l_{\mu_0}(u, y_1^1) > C_\star$ and u lies on μ_0 between y_1 and y_1^1 .

Claim. For every decomposition $p \equiv p_1 p_2$, where p_1 is a proper initial edge-subpath of p , cancellations in the product $p_2 q \cdot [f(p_1)]$ don't touch p_2 .

Indeed, since p_1 is an initial subpath of μ_0 , by Cooper's lemma, we have $[f(p_1)] = ab$ for some initial subpath a of $[f(\mu_0)]$ and some path b of length at most $C_\star$. This and the fact that $l_{\mu_0}(u, y_1^1) > C_\star$ imply the claim.

Let $\mu_1 \equiv \hat{f}^{l(p)}(\mu_0)$. Clearly, μ_1 lies in the μ_0 -subgraph. We assert that $N_r(\mu_1) < N_r(\mu_0)$. To prove this, we describe the path μ_1 precisely. First, by the above claim we have $\mu_1 \equiv [qf(p)]$. Then $\mu_1 \equiv \mu_{uy_1^1} \cdot \ell$, where ℓ is the subpath of $[f(\mu_0)]$ from $y_1^1 = z^0$ to $f(u)$. Since ℓ is r -legal, the r -cancellation points of μ_1 are contained in the set $\{y_1, \dots, y_k, y_1^1\} \setminus \{y_1\}$. However y_1^1 is not an r -cancellation point of μ_1 by Section 5. So, $N_r(\mu_1) < N_r(\mu_0)$.




Figure 17.

The path μ_1 is depicted by the bold line.

Case 2.2. Suppose that $l_{\mu_0}(y_1, y_1^1) \leq R_\star + C_\star$.

Let p be the initial subpath of μ_0 which terminates at y_1 (see Figure 18) and $\mu_0 \equiv pq$ for some q . We set $\mu_1 = \widehat{f}^{l(p)}(\mu_0)$. As in Case 1, one can prove that $\mu_1 \equiv [qf(p)]$ and hence $\mu_1 \equiv [\mu_{y_1 y_1^1} \cdot \ell]$, where $\mu_{y_1 y_1^1}$ is the initial subpath of q from y_1 to y_1^1 and ℓ is the final subpath of $[f(p)]$ from y_1^1 to $f(y_1)$. Using Cooper's lemma, we deduce that the l -length of ℓ is at most $C_\star$, hence

$$l(\mu_1) \leq l_{\mu_0}(y_1, y_1^1) + l_{[f(p)]}(y_1^1, f(y_1)) \leq R_\star + 2C_\star.$$

This contradicts the condition iii).




Figure 18.

The path μ_1 is depicted by the bold line.

Case 3. Suppose that both, y_k and y_1^1 don't lie in I_0 .

Then $y_1, \dots, y_k, y_1^1, \dots, y_k^1$ don't lie in I_0 .

We may assume that z^0 is an r -cancellation point in $[\mu_0 f(\mu_0)]$.

Indeed, suppose that this is not valid. We set $\mu_1 = \widehat{f}^{l(p)}(\mu_0)$, where p is the initial subsegment of μ_0 with the endpoint y_1 (see Figure 19). Then μ_1 lies in the μ_0 -subgraph and the r -cancellation points of μ_1 are contained in the set $\{y_2, \dots, y_k, z^0\}$. But z^0 is not an r -cancellation point in μ_1 , since we have supposed that z^0 is not

an r -cancelation point in $[\mu_0 f(\mu_0)]$. So, $N(\mu_1) = N(\mu_0) - 1$ and we can proceed by induction.




Figure 19.

The path μ_1 is depicted by the bold line.

Thus, we may assume that $y_1, \dots, y_k, z^0, y_1^1, \dots, y_k^1$ are all r -cancelation points in $[\mu_0 f(\mu_0)]$. Since μ_0 is r -superstable, they are non-deletable there.

Let $A(y_1^1)$ be the r -cancelation area of y_1^1 in $[f(\mu_0)]$. Let $A'(z^0)$ and $A'(y_1^1)$ be the r -cancelation areas of z^0 and y_1^1 in $[\mu_0 f(\mu_0)]$, respectively.

Claim. We have $A(y_1^1) = A'(y_1^1)$. In particular, $\alpha(A(y_1^1))$ lies to the right from z^0 in $[\mu_0 f(\mu_0)]$.

Proof. Since the interiors of $A'(z^0)$ and $A'(y_1^1)$ don't intersect, $A'(y_1^1)$ lies to the right from z^0 in $[\mu_0 f(\mu_0)]$. Hence $A'(y_1^1)$ lies in $[f(\mu_0)]$. Therefore $A(y_1^1) = A'(y_1^1)$.




Figure 20.

The path μ_1 is depicted by the bold line.

Let u be the initial vertex of the r -cancelation area $A(y_1)$. The point u divides μ_0 into two subpaths which we denote by p and q , so $\mu_0 \equiv pq$. Now we set

$\mu_1 \equiv \widehat{f}^{l(p)}(\mu_0)$. Clearly, μ_1 lies in the μ_0 -subgraph. As in Case 1, we can prove that $\mu_1 \equiv [qf(p)]$ and hence μ_1 is the subpath of $[\mu_0 f(\mu_0)]$ from $\alpha(A(y_1))$ to $\alpha(A(y_1^1))$. We prove that μ_1 is A -perfect. Obviously, the A -decomposition of μ_1 starts on the r -cancelation area $A(y_1)$. It remains to prove that the turn in $\mu_1 \cdot [f(\mu_1)]$ between μ_1 and $[f(\mu_1)]$ is legal. The first edge of $[f(\mu_1)]$ is the first edge of $[f(A(y_1))] = A(y_1^1)$, so this edge lies in H_r . If the last edge of μ_1 lies in G_{r-1} , then the turn is mixed and so legal. If the last edge of μ_1 lies in H_r , then this turn is an r -turn. But this turn lies in the r -legal subpath of $[\mu_0 f(\mu_0)]$ from z^0 to y_1^1 , hence it is r -legal.

Thus, μ_1 is A -perfect and lies in $\widetilde{M}$; the proof is completed. $\square$

Proposition 13.4. *Suppose that μ is a reduced f -path in G_r and that the μ -subgraph M is infinite. Then we can algorithmically find a vertex μ' in M with one of the following properties:*

- 1) *the f -path μ' lies in G_{r-1} ,*
- 2) *μ' is r -perfect,*
- 3) *μ' is A -perfect.*

Proof. Let $D_f(\mu)$ be the component of D_f containing μ . We claim that there exists a vertex $\widehat{\mu}$ in the μ -subgraph, which satisfies conditions ii) and iii) of Proposition 13.3. This follows from the facts that $\pi_1(D_f(\mu), \mu)$ is finitely generated (see Lemma 5.3) and that the number of dead and repelling vertices in D_f is finite. By Proposition 13.3, where we replace μ by $\widehat{\mu}$, there exists a vertex μ' in the $\widehat{\mu}$ -subgraph which satisfies one of the properties 1)-3). Clearly, μ' lies in the μ -subgraph and we can find μ' just by checking consecutive vertices in the μ -subgraph. $\square$

14. E -PERFECT PATHS

In this section we will use that the subdivided relative train track $f : \Gamma \rightarrow \Gamma$ satisfies the condition (Pol) in Theorem 4.5 (see Proposition 9.5).

Let H_r be a polynomial stratum, let E be the unique (up to inversion) edge of H_r , and let $f(E) \equiv E \cdot \sigma$, where σ is a path in G_{r-1} .

For an edge-path $\mu \subset G_r$, let $\mathcal{N}(\mu)$ be the number of edges from H_r in μ .

We introduce some notations: For every $Q \in \{E, \overline{E}\}$ and every $i \geq 1$, we have $f^i(Q) \equiv c_{i,Q} Q d_{i,Q}$, where

$$c_{i,Q} = \begin{cases} 1_{\alpha(Q)} & \text{if } Q = E, \\ f^{i-1}(\bar{\sigma}) \dots f(\bar{\sigma})\bar{\sigma} & \text{if } Q = \overline{E}, \end{cases}$$

$$d_{i,Q} = \begin{cases} 1_{\omega(Q)} & \text{if } Q = \overline{E}, \\ \sigma f(\sigma) \dots f^{i-1}(\sigma) & \text{if } Q = E. \end{cases}$$

We set $c_{0,Q} = \mathbf{1}_{\alpha(Q)}$ and $d_{0,Q} = \mathbf{1}_{\omega(Q)}$. Then, for every $i \geq 0$, we have

$$f^i(Q) \equiv c_{i,Q} Q d_{i,Q}.$$

Definition 14.1. The vertex $\mu \in D_f$ is called *E-perfect* if $\mu \equiv E_1 b_1 E_2 \dots E_k b_k$, where $k \geq 1$, $E_1, \dots, E_k \in \{E, \overline{E}\}$ and $b_1, \dots, b_k$ are paths in G_{r-1} , and $\mathcal{N}(\mu') = \mathcal{N}(\mu)$ for every vertex μ' in the μ -subgraph.

Proposition 14.2. Let H_r be an exponential stratum with a unique (up to inversion) edge E . Let $\mu \equiv E_1 b_1 \dots E_k b_k$ be a reduced f -path in G_r , where $E_1, \dots, E_k \in \{E, \overline{E}\}$ and $b_1, \dots, b_k \subset G_{r-1}$. For $1 \leq j \leq k$ and $i \geq 1$, we set

$$\mu_{0,j} \equiv [E_j b_j \dots E_k b_k f(E_1 b_1 \dots E_{j-1} b_{j-1})],$$

$$\mu_{i,j} \equiv [\overline{c_{i,E_j}} f^i(\mu_{0,j}) f(c_{i,E_j})].$$

The elements of the sequence $\mu_{0,1}, \dots, \mu_{0,k}, \mu_{1,1}, \dots, \mu_{1,k}, \dots$ will be denoted by $\mu_1, \mu_2, \dots$. (Clearly, all these elements are reduced f -paths in G_r .) Then the following statements are satisfied.

- (1) If $\mathcal{N}(\mu_i) = k$ for $1 \leq i \leq k$, then $\mathcal{N}(\mu_i) = k$ for all $i \geq 1$. In this case μ is *E-perfect* and all μ_i lie in the μ -subgraph. Moreover, μ_{i+1} is the first vertex of the μ -subgraph after μ_i which, considered as an f -path, begins on E or $\overline{E}$.
- (2) If for some $1 \leq s < k$ we have $\mathcal{N}(\mu_1) = \dots = \mathcal{N}(\mu_s) = k$, but $\mathcal{N}(\mu_{s+1}) < k$, then $\mu_1, \dots, \mu_s$, and μ_{s+1} lie in the μ -subgraph. In this case μ is not *E-perfect*.
In particular, we can find a vertex in the μ -subgraph which is *E-perfect* or lies in G_{r-1} (considered as an f -path).
- (3) If μ is *E-perfect*, then a vertex in the μ -subgraph is *E-perfect* if and only if it, considered as an f -path, begins on E or $\overline{E}$.

Proof. To prove the first claim, it suffices to verify that $\mathcal{N}(\mu_{0,j}) = \mathcal{N}(\mu_{i,j})$ for all $1 \leq j \leq k$ and $i \geq 1$. This is equivalent to $\mathcal{N}(\mu_{0,j}) = \mathcal{N}([f^i(\mu_{0,j})])$.

For $k = 1$ this is evident. Suppose that $k \geq 2$. By assumption we have $\mathcal{N}(\mu_{0,j}) = k$, hence $\mu_{0,j} \equiv Q_1 \cdot c_1 \cdot \dots \cdot Q_k \cdot c_k$, where $Q_1, \dots, Q_k \in \{E, \overline{E}\}$ and $c_1, \dots, c_k$ are paths in G_{r-1} . Then

$$[f^i(\mu_{0,j})] \equiv [c_{i,Q_1} Q_1 d_{i,Q_1} \cdot f^i(c_1) \cdot \dots \cdot c_{i,Q_k} Q_k d_{i,Q_k} \cdot f^i(c_k)].$$

Consider a subword $Q_s c_s Q_{s+1}$ of $\mu_{0,j}$ and the corresponding subword of $f^i(\mu_{0,j})$:

$$Q_s d_{i,Q_s} f^i(c_s) c_{i,Q_{s+1}} Q_{s+1}.$$

We prove that the reduced form of the corresponding subword contains Q_s and Q_{s+1} . Suppose the contrary. Then Q_s is inverse to Q_{s+1} . Since $Q_s c_s Q_{s+1}$ is reduced, c_s is a nontrivial loop in G_{r-1} .

Case 1. Suppose that $Q_s = E$ and $Q_{s+1} = \overline{E}$.

Then $Q_s d_{i,Q_s} f^i(c_s) c_{i,Q_{s+1}} Q_{s+1} \equiv E \sigma f(\sigma) \dots f^{i-1}(\sigma) f^i(c_s) f^{i-1}(\bar{\sigma}) \dots f(\bar{\sigma}) \bar{\sigma} \overline{E}$.

Since c_s is a nontrivial loop, the loop between E and $\overline{E}$ is this path is nontrivial as well. Hence the reduced form of this path contains E and $\overline{E}$.

Case 2. Suppose that $Q_s = \overline{E}$ and $Q_{s+1} = E$.

Then $Q_s d_{i,Q_s} f^i(c_s) c_{i,Q_{s+1}} Q_{s+1} \equiv f^{i-1}(\bar{\sigma}) \dots f(\bar{\sigma}) \bar{\sigma} \overline{E} c_s E \sigma f(\sigma) \dots f^{i-1}(\sigma)$.

Clearly, the reduced form of this path contains $\overline{E}$ and E too.

The other claims of this proposition can be routinely verified. $\square$

Proposition 14.3. *For every two vertices μ, τ in D_f , where μ is E -perfect and the μ -subgraph is infinite, one can algorithmically decide, whether τ lies in the μ -subgraph.*

Proof. We may assume that the f -path τ lies in G_r . Moving along the τ -subgraph, we will find a vertex which is E -perfect or lies in G_{r-1} (considered as an f -path). Hence, without loss of generality, we may assume that τ is E -perfect. By Proposition 14.2, τ lies in the μ -subgraph if and only if $\tau \equiv \mu_{i,j}$ for some $i \geq 0$ and $1 \leq j \leq k$. One can easily check that

$$c_{i,E_j} = \begin{cases} \mathbf{1}_{\alpha(E)} & \text{if } E_j = E, \\ \overline{f^i(E)} E & \text{if } E_j = \overline{E}, \end{cases}$$

Using this, we deduce

$$\mu_{i,j} \equiv \begin{cases} [f^i(\mu_{0,j})] & \text{if } E_j = E, \\ [\overline{E} f^i(E) \cdot f^i(\mu_{0,j}) \cdot f(\overline{f^i(E)} E)] & \text{if } E_j = \overline{E}. \end{cases}$$

Simplifying, we have

$$\mu_{i,j} \equiv \begin{cases} [f^i(\mu_{0,j})] & \text{if } E_j = E, \\ [\overline{E} \cdot f^i(E \mu_{0,j} \overline{f(E)}) \cdot f(E)] & \text{if } E_j = \overline{E}. \end{cases} \quad (14.1)$$

Thus, $\tau \equiv \mu_{i,j}$ if and only if

$$\begin{cases} \tau \equiv [f^i(\mu_{0,j})] & \text{if } E_j = E, \\ [E \tau \overline{f(E)}] = [f^i(E \mu_{0,j} \overline{f(E)})] & \text{if } E_j = \overline{E}. \end{cases}$$

By Corollary 4.8, we can algorithmically decide the question about the existence of such i, j . $\square$

15. FINITENESS AND MEMBERSHIP PROBLEMS FOR μ -SUBGRAPHS

In this section we will use that the subdivided relative train track $f : \Gamma \rightarrow \Gamma$ satisfies the condition (Pol) in Theorem 4.5 (see Proposition 9.5). For such f we prove Propositions 15.2 and 15.5 which solve Problems 2 and 1 from Section 4. This will complete the proof of the main Theorem 1.1.

Lemma 15.1. *Let H_r be an exponential stratum with the Perron-Frobenius value λ_r and let $\mu \subset G_r$ be an r -perfect path with $l(\mu) \geq K$, where K is the constant from Proposition 9.5. 8). Let $\mu = \mu_1, \mu_2, \dots$ be consecutive vertices of the μ -subgraph in D_f . Then, for all $i \geq 1$, we have*

$$L_r(\mu_{i+1}) \geq L_r(\mu_i) > 0$$

and

$$L_r(\mu_{m_i}) = \lambda_r^i L_r(\mu),$$

where $m_1 < m_2 < \dots$ are computable numbers from Proposition 10.2.

Proof. By Proposition 9.5. 8), there exists $k \geq 1$ such that μ_k lies in G_{r-1} or satisfies $L_r(\mu_k) > 0$. By the claim (3) of Proposition 10.2, μ_k contains edges from H_r , hence $L_r(\mu_k) > 0$. This implies $L_r(\mu_1) > 0$. Other formulas follow from the claims (3) and (2) of Proposition 10.2. $\square$

Proposition 15.2. *Given a vertex μ in D_f , one can decide whether the μ -subgraph is finite or not.*

Proof. Let r be the minimal number such that the f -path μ lies in G_r .

First suppose that H_r is an exponential stratum. We claim that there exists $n \geq 0$ such that the vertex $\mu' = \widehat{f}^n(\mu)$ of the μ -subgraph satisfies one of the following properties:

- 1) the f -path μ' lies in G_{r-1} ,
- 2) μ' is r -perfect,
- 3) μ' is A -perfect,
- 4) the vertices $\mu, \widehat{f}(\mu), \dots, \widehat{f}^{n-1}(\mu)$ are different and μ' coincides with one of them,
- 5) μ' is a dead vertex.

Indeed, if the μ -subgraph is infinite, then, by Proposition 13.4, there exists n such that $\mu' = \widehat{f}^n(\mu)$ satisfies one of the properties 1)-3). If the μ -subgraph is finite, there obviously exists n such that $\mu' = \widehat{f}^n(\mu)$ satisfies one of the properties 4)-5).

Observe that for every $n \geq 0$, we can algorithmically verify whether $\mu' = \widehat{f}^n(\mu)$ satisfies one of the conditions 1)-5).

So, starting from μ and following along preferable directions, we can find $n \geq 0$ such that the corresponding μ' satisfies one of the above conditions. If 4) or 5) is satisfied, then the μ -subgraph is finite.

If 1) is satisfied, we can apply induction.

If 2) is satisfied, then the μ -subgraph is infinite by Lemma 15.1.

Suppose that 3) is satisfied. By Proposition 10.4 (2), there exist natural numbers $m_{1,1} < m_{2,1} < m_{3,1} < \dots$ such that $\widehat{f}^{m_{i,1}}(\mu') \equiv [f^i(\mu')]$, $i > 0$. Hence, the μ' -subgraph is finite if and only if there exist $0 < i < j$ such that $[f^i(\mu')] = [f^j(\mu')]$.

This problem is decidable by Corollary 4.9.

Now suppose that H_r is a polynomial stratum. By Proposition 14.2, we can find a vertex μ' in the μ -subgraph with one of the following properties:

- 1) the f -path μ' lies in G_{r-1} ,
- 2) μ' is E -perfect.

In the first case we can apply induction. Suppose that μ' is E -perfect. By Proposition 14.2, the μ' -subgraph contains the vertices $\mu'_{i,1}$, $i \geq 0$. Hence, the μ' -subgraph is finite if and only if there exist $0 \leq i < l$ such that $\mu'_{i,1} = \mu'_{l,1}$. (Recall that $\mu'_{0,1} = \mu'$ and E_1 is the first edge of μ' .) By the formula (14.1) this is equivalent to the equality:

$$\begin{cases} [f^i(\mu')] = [f^l(\mu')] & \text{if } E_1 = E, \\ [f^i(E\mu'f(E))] = [f^l(E\mu'f(E))] & \text{if } E_1 = \overline{E}. \end{cases}$$

This problem is decidable by Corollary 4.9.

If H_r is a zero stratum, then following along the μ -subgraph at most $l(\mu)$ steps, we can find a vertex μ' in the μ -subgraph such that the f -path μ' lies in G_{r-1} . Then we can apply induction. $\square$

Proposition 15.3. *For every two vertices μ, τ in D_f , where μ is r -perfect, one can decide whether τ lies in the μ -subgraph.*

Proof. We may assume that τ is an f -path in G_r . Let $\mu = \mu_0, \mu_1, \dots$, be consecutive vertices of the μ -subgraph. By Lemma 15.1, we compute the minimal i such that $L_r(\mu_i) > L_r(\tau)$. Then it suffices to compare τ with the vertices $\mu_0, \mu_1, \dots, \mu_{i-1}$. $\square$

Proposition 15.4. *For every two vertices μ, τ in D_f , where μ is A -perfect, one can decide whether τ lies in the μ -subgraph.*

Proof. Due to Proposition 15.2, we may assume that the μ -subgraph is infinite. Let $\mu \equiv A_1 b_1 \dots A_k b_k$ be the A -decomposition of μ . We use the following notation from Proposition 10.4:

$$\begin{aligned} \mu_{0,j} &\equiv [A_j b_j \dots A_k b_k f(A_1 b_1 \dots A_{j-1} b_{j-1})], \\ \mu_{i,j} &\equiv [f^i(\mu_{0,j})], \end{aligned}$$

where $1 \leq j \leq k$ and $i \geq 1$. By the statement (4) of Proposition 10.4 the following holds: for every vertex σ in the μ -subgraph, at least one of the paths $\sigma, \widehat{f}(\sigma), \dots, \widehat{f}^{l(\sigma)}(\sigma)$ coincides with $\mu_{i,j}$ for some i, j .

Thus, we first decide, whether one of the vertices $\tau, \widehat{f}(\tau), \dots, \widehat{f}^{l(\tau)}(\tau)$ coincides with $\mu_{i,j}$ for some i, j . This can be done with the help of Corollary 4.8.

If the answer is negative, then τ does not lie in the μ -subgraph.

If it is positive, then we can find t, i, j such that $\widehat{f}^t(\tau) \equiv \mu_{i,j}$. Recall that by the statement (2) of Proposition 10.4, $\mu_{i,j} \equiv \widehat{f}^{m_{i,j}}(\mu)$ for some computable $m_{i,j}$. Then τ lies in the μ -subgraph if and only if $m_{i,j} \geq t$ and $\tau \equiv \widehat{f}^{m_{i,j}-t}(\mu)$. $\square$

Proposition 15.5. *Given two vertices μ, τ in D_f , one can decide whether τ lies in the μ -subgraph.*

Proof. By Proposition 15.2, we can decide whether these subgraphs are finite or not. If the μ -subgraph is finite, we can list all its vertices and verify, whether τ is one of them. Suppose that the μ -subgraph is infinite. Then, if the τ -subgraph is finite, the vertex τ cannot lie in the μ -subgraph. So, we can assume that the τ -subgraph is also infinite.

Let r be the minimal number such that μ lies in G_r . We will use induction on r .

First suppose that H_r is an exponential stratum. Then, by Proposition 13.4, we can find a vertex μ' in the μ -subgraph with one of the following properties:

- 1) the f -path μ' lies in G_{r-1} ,
- 2) μ' is r -perfect,
- 3) μ' is A -perfect,

First we check, whether τ belongs to the segment of the μ -subgraph from μ to μ' . If yes, we are done. If not, we replace μ by μ' and consider the above cases.

In Case 1) we can proceed by induction, in Case 2) by Proposition 15.3 and in Case 3) by Proposition 15.4.

Now suppose that H_r is a polynomial stratum. Then, by Proposition 14.2, we can find a vertex μ' in the μ -subgraph with one of the following properties:

- 1) the f -path μ' lies in G_{r-1} ,
- 2) μ' is E -perfect.

In the first case we can apply induction and in the second case Proposition 14.3.

Finally, if H_r is a zero stratum, we can follow along the μ -subgraph until we arrive at a vertex $\mu' \in D_f$ which, considered as an f -path, lies in G_{r-1} . Then we can apply induction. $\square$

REFERENCES

- [1] G. Baumslag, A. G. Myasnikov, V. Shpilrain, *Open problems in combinatorial group theory*. Second edition, (Contemp. Math., **296**), Providence, RI, Am. Math. Soc., 2002.
- [2] G. M. Bergman, *Supports of derivations, free factorizations, and ranks of fixed subgroups in free groups*, Trans. Am. Math. Soc. **351**, N 4 (1999), 1531–1550.
- [3] M. Bestvina, *$\mathbb{R}$ -trees in topology, geometry, and group theory*, Handbook of geometric topology, 5591, North-Holland, Amsterdam, 2002.
- [4] M. Bestvina, M. Feighn, M. Handel, *The Tits alternative for $\text{Out}(F_n)$ I: Dynamics of exponentially-growing automorphisms*, Ann. Math., **151**, N 2 (2000), 517–623.
- [5] M. Bestvina, M. Feighn, M. Handel, *The Tits alternative for $\text{Out}(F_n)$ II: A Kolchin type theorem*, Ann. Math., **161**, N 1 (2005) Pages 1–59.
- [6] M. Bestvina, M. Feighn, M. Handel, *Solvable subgroups of $\text{Out}(F_n)$ are virtually abelian*, Geometriae Dedicata, **104**, N 1 (2004), 71–96.
- [7] M. Bestvina, M. Handel, *Train tracks and automorphisms of free groups*, Ann. Math. (2), **135**, N 1 (1992), 1–53.
- [8] O. Bogopolski, *Introduction to group theory*, EMS publishing house, 2008 (see Section 3 there).
- [9] O. Bogopolski, *Classification of automorphisms of the free group of rank 2 by ranks of fixed-point subgroups*, J. Group Theory, **3**, N 3 (2000), 339–351.
- [10] O. Bogopolski, A. Martino, O. Maslakova, E. Ventura, *Free-by-cyclic groups have solvable conjugacy problem*, Bulletin of the London Math. Soc., **38**, part 5 (2006), 787–794.
- [11] M.R. Bridson, D. Groves, *The quadratic isoperimetric inequality for mapping tori of free group automorphisms*, Mem. Am. Math. Soc. 955, xii, 152 pp. (2010).
- [12] P. Brinkmann, *Hyperbolic automorphisms of free groups*, Geom. Funct. Anal. **10**, N 5 (2000), 1071–1089.
- [13] P. Brinkmann, *Detecting automorphic orbits in free groups*, Journal of Algebra, **324** (2010), 1083–1097.
- [14] P. Brinkmann, *Dynamics of free groups automorphisms*, preprint, (2003), 40 pp.
- [15] D.J. Collins, E.C. Turner, *All automorphisms of free groups with maximal rank fixed subgroups*, Math. Proc. Camb. Philos. Soc. **119**, N 4 (1996), 615–630.
- [16] M. M. Cohen, M. Lustig, *On the dynamics and the fixed subgroup of a free group automorphism*, Invent. Math., **96**, N 3 (1989), 613–638.
- [17] D. Cooper, *Automorphisms of free groups have finitely generated fixed point sets*, J. Algebra, **111**, N 2 (1987), 453–456.
- [18] W. Dicks, E. Ventura, *The group fixed by a family of injective endomorphisms of a free group* (Contemp. Math., **195**), Providence, RI, Am. Math. Soc., 1996.
- [19] J. L. Dyer, G. P. Scott, *Periodic automorphisms of free groups*, Commun. Algebra, **3**, N 3 (1975), 195–201.
- [20] M. Feighn, M. Handel, *Abelian subgroups of $\text{Out}(F_n)$* , Geometry and Topology, **13** (2009), 1657–1727.
- [21] D. Gaboriau, A. Jaeger, G. Levitt, M. Lustig, *An index for counting fixed points of automorphisms of free groups*, Duke Math. J., **93**, N 3 (1998), 425–452.
- [22] D. Gaboriau, G. Levitt, *The rank of actions on R -trees*, Ann. Sci. Ecole Norm. Sup. **28** (1995), 549–570.
- [23] D. Gaboriau, G. Levitt, M. Lustig, *A dendrological proof of the Scott conjecture for automorphisms of free groups*, Proc. Edinburgh Math. Soc. (2), **41**, N 2 (1998), 325–332.

- [24] S. M. Gersten, *On fixed points of automorphisms of finitely generated free groups*, Bull. Amer. Math. Soc., New Ser. **8** (1983), 451–454.
- [25] S. M. Gersten, *Fixed points of automorphisms of free groups*, Adv. in Math., **64**, N 1 (1987), 51–85.
- [26] R. Z. Goldstein, E. C. Turner, *Fixed subgroups of homomorphisms of free groups*, Bull. London Math. Soc., **18** (1986), 468–470.
- [27] R. Z. Goldstein, E. C. Turner, *Automorphisms of free groups and their fixed points*, Inv. Math., **78**, N 1 (1984), 1–12.
- [28] W. Imrich, E. C. Turner, *Endomorphisms of free groups and their fixed points*, Math. Proc. Cambridge Philos. Soc., **105** (1989), 421–422.
- [29] S. Kalajdzevski, *Automorphism group of a free group: Centralizers and stabilizers*, J. Algebra, **150**, N 2 (1992), 435–502.
- [30] O. S. Maslakova, *Fixed point subgroup of an automorphism of a free group*, Algebra and Logic, **42**, N 4 (2003), 422–472 (in Russian).
- [31] O. S. Maslakova, *Fixed point subgroup of an automorphism of a free group*, Ph.D Thesis (2004) (in Russian).
- [32] F. Paulin, *Sur les automorphismes extérieurs des groupes hyperboliques*, Ann. Inst. Fourier **39** (1989), 651–662.
- [33] Z. Sela, *The Nielsen-Thurston classification and automorphisms of a free group I*, Duke Math. J., **84**, N 2 (1996), 379–397.
- [34] J.R. Stallings, *Topology of finite graphs*, Inv. Math., **71**, N 3 (1983), 551–566.
- [35] J.R. Stallings, *Graphical theory of automorphisms of free groups*, Ann. Math. Stud. **111** (1987), 79–106.
- [36] S. Thomas, *Fixed points of automorphisms of finitely generated free groups*, Proc. Amer. Math. Soc., **103** (1988), 333.
- [37] E. C. Turner, *Finding invisible Nielsen paths for a train tracks map*, Proc. of a workshop held at Heriot-Watt Univ., Edinburg, 1993 (Lond. Math. Soc. Lect. Note Ser., **204**), Cambridge, Cambridge Univ. Press., 1995, 300–313.